

2022–2024

arXiv:2411.02368v4 [quant-ph] 5 Jun 2025

Optimal estimates of trace distance between bosonic Gaussian states and applications to learning

Lennart Bittel[1], Francesco Anna Mele[2], Antonio Anna Mele[1], Salvatore Tirone[3,4], and Ludovico Lami[5,3,4,6]

[1]Dahlem Center for Complex Quantum Systems, Freie Universität Berlin, 14195 Berlin, Germany
[2]NEST, Scuola Normale Superiore and Istituto Nanoscienze, Piazza dei Cavalieri 7, IT-56126 Pisa, Italy
[3]QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands
[4]Korteweg–de Vries Institute for Mathematics, University of Amsterdam, Science Park 105-107, 1098 XG Amsterdam, the Netherlands
[5]Scuola Normale Superiore, Piazza dei Cavalieri 7, 56126 Pisa, Italy
[6]Institute for Theoretical Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands

Gaussian states of bosonic quantum systems enjoy several technological applications and are ubiquitous in nature. Their significance lies in their simplicity, which in turn rests on the fact that they are uniquely determined by two experimentally accessible quantities, their first and second moments. But what if these moments are only known *approximately*, as is inevitable in any realistic experiment? What is the resulting error on the Gaussian state itself, as measured by the most operationally meaningful metric for distinguishing quantum states, namely, the trace distance? In this work, we fully resolve this question by demonstrating that if the first and second moments are known up to an error ε, the trace distance error on the state also scales as ε, and this functional dependence is optimal. To prove this, we establish tight bounds on the trace distance between two Gaussian states in terms of the norm distance of their first and second moments. As an application, we improve existing bounds on the sample complexity of tomography of Gaussian states. In our analysis, we introduce the general notion of derivative of a Gaussian state and uncover its fundamental properties, enhancing our understanding of the structure of the set of Gaussian states.

Introduction.— Continuous variable (CV) systems [1?–5], such as quantum optical systems, play a key role in our quantum technologies, including quantum computation [6–14], communication [15–22], and sensing [23–27]. Additionally, CV systems have become increasingly popular in efforts to demonstrate quantum advantage, particularly through boson sampling [28–30] and quantum simulation experiments [31]. Among all CV quantum states, *Gaussian states* [2] are arguably among the most important ones. Indeed, on the one hand Gaussian states are the most common class of states arising in Nature, while on the other they are relatively simple to analyse mathematically.

Gaussian states are the quantum analog of Gaussian probability distributions. Just as a Gaussian probability distribution is uniquely identified by its classical first moment and its classical covariance matrix, a Gaussian state is uniquely identified by its *first moment* $\mathbf{m}$ and its *covariance matrix* V [2]. Importantly, these quantities can be experimentally estimated through methods such as homodyne or heterodyne detection [2], which are routinely performed in quantum optical laboratories. Thus, estimating a Gaussian state requires only the estimation of its first moment and covariance matrix. In practice, however, we can never estimate the first moment and the covariance matrix *exactly*; instead, we will always have a non-zero error ε in such estimates, meaning that we can only estimate a Gaussian state up to some error. The goal of this work is to understand how an estimation error ε in the first moment and the covariance matrix of a Gaussian state propagates into the overall error in estimating the Gaussian state itself:

Problem 1 (Informal version)**.** *Let $\rho(V,\mathbf{m})$ be the Gaussian state with first moment $\mathbf{m}$ and covariance matrix V. Let $\tilde{V}$ and $\tilde{\mathbf{m}}$ be ε-approximations of V and $\mathbf{m}$, respectively:*

$$\tilde{\mathbf{m}} \approx_\varepsilon \mathbf{m}\,, \qquad \tilde{V} \approx_\varepsilon V\,. \tag{1}$$

The Gaussian state $\rho(\tilde{V},\tilde{\mathbf{m}})$ with first moment $\tilde{\mathbf{m}}$ and covariance matrix $\tilde{V}$ constitutes an approximation of $\rho(V,\mathbf{m})$:

$$\rho(\tilde{V},\tilde{\mathbf{m}}) \approx \rho(V,\mathbf{m})\,. \tag{2}$$

The problem is to understand what is the error incurred in the approximation in Eq. (2) in terms of the error ε incurred in the approximations of the first moment and of the covariance matrix.

Lennart Bittel: bittel@hhu.de
Francesco Anna Mele: francesco.mele@sns.it
Antonio Anna Mele: a.mele@fu-berlin.de
Salvatore Tirone: s.tirone@uva.nl
Ludovico Lami: ludovico.lami@gmail.com