



SciPost Physics Submission

Parallel assembly of neutral atom arrays with an SLM using linear phase interpolation

Ivo H. A. Knottnerus[1,2,3], Yu Chih Tseng (曾于誌)[1,2], Alexander Urech[1,2], Robert J. C. Spreeuw[1,2] and Florian Schreck[1,2⋆]

1 Van der Waals-Zeeman Institute, Institute of Physics, University of Amsterdam, Science Park 904, 1098XH Amsterdam, The Netherlands
2 QuSoft, Science Park 123, 1098XG Amsterdam, The Netherlands
3 Eindhoven University of Technology, P.O. Box 513, 5600MB Eindhoven, The Netherlands

⋆ slmsorting@strontiumbec.com

arXiv:2501.01391v4 [quant-ph] 23 Aug 2025

Abstract

We present fast parallel rearrangement of single atoms in optical tweezers into arbitrary geometries by updating holograms displayed by an ultra fast spatial light modulator. Using linear interpolation of the tweezer position and the optical phase between the start and end arrays, we can calculate and display holograms every few ms, limited by technology. To show the versatility of our method, we sort the same atomic sample into multiple geometries with success probabilities of 0.996(2) per rearrangement cycle. This makes the method a useful tool for rearranging large atom arrays for quantum computation and quantum simulation.

Copyright attribution to authors.
This work is a submission to SciPost Physics.
License information to appear upon publication.
Publication information to appear upon publication.

Received Date
Accepted Date
Published Date

Contents

1 Introduction 2
2 Parallel rearrangement of atoms with SLM 3
3 Linear phase interpolation for flicker-free transport 5
4 Scaling to large atom arrays 8
5 Discussion and Outlook 10
6 Conclusion 11
A Experimental details 11
B Data analysis and fidelities 12
C Derivation of phase slip per displacement 14
References 15

