


Trotter error and gate complexity of the SYK and sparse SYK models

Yiyuan Chen[1], Jonas Helsen[2,3], and Maris Ozols[4,1,2]

[1]Korteweg-de Vries Institute for Mathematics, University of Amsterdam, The Netherlands
[2]QuSoft, Amsterdam, The Netherlands
[3]CWI, Amsterdam, The Netherlands
[4]Institute for Logic, Language, and Computation, University of Amsterdam, The Netherlands

arXiv:2502.18420v1 [quant-ph] 25 Feb 2025

The Sachdev–Ye–Kitaev (SYK) model is a prominent model of strongly interacting fermions that serves as a toy model of quantum gravity and black hole physics. In this work, we study the Trotter error and gate complexity of the quantum simulation of the SYK model using Lie–Trotter–Suzuki formulas. Building on recent results by Chen and Brandão [CB24] – in particular their uniform smoothing technique for random matrix polynomials –, we derive bounds on the first- and higher-order Trotter error of the SYK model, and subsequently find near-optimal gate complexities for simulating these models using Lie–Trotter–Suzuki formulas. For the k-local SYK model on n Majorana fermions, our gate complexity estimates for the first-order Lie–Trotter–Suzuki formula scales with $\mathcal{O}(n^{k+\frac{5}{2}}t^2)$ for even k and $\mathcal{O}(n^{k+3}t^2)$ for odd k, and the gate complexity of simulations using higher-order formulas scales with $\mathcal{O}(n^{k+\frac{1}{2}}t)$ for even k and $\mathcal{O}(n^{k+1}t)$ for odd k. Given that the SYK model has $\Theta(n^k)$ terms, these estimates are close to optimal. These gate complexities can be further improved upon in the context of simulating the time-evolution of an arbitrary fixed input state $|\psi\rangle$, leading to a $\mathcal{O}(n^2)$-reduction in gate complexity for first-order formulas and $\mathcal{O}(\sqrt{n})$-reduction for higher-order formulas.

We also apply our techniques to the sparse SYK model, which is a simplified variant of the SYK model obtained by deleting all but a $\Theta(n)$ fraction of the terms in a uniformly i.i.d. manner. We compute the average (over the random term removal) gate complexity for simulating this model using higher-order formulas to be $\mathcal{O}(n^2t)$, a bound that also holds for a general class of sparse Gaussian random Hamiltonians. Similar to the full SYK model, we obtain a $\mathcal{O}(\sqrt{n})$-reduction simulating the time-evolution of an arbitrary fixed input state $|\psi\rangle$.

Our results highlight the potential of Lie–Trotter–Suzuki formulas for efficiently simulating the SYK and sparse SYK models, and our analytical methods can be naturally extended to other Gaussian random Hamiltonians.

Contents

1 Introduction **2**
1.1 Previous work . . . 3
1.2 Summary of results for the SYK model . . . 4
1.2.1 Error bounds . . . 4
1.2.2 Gate complexity . . . 5
1.3 Summary of results for the sparse SYK model . . . 6
1.4 Comparison with numerical simulations . . . 7
1.5 Structure of the paper . . . 8
1.6 Acknowledgements . . . 8

2 Preliminaries **9**

