

Governing fields for hyperelliptic function fields

Joppe Stokvis *

May 27, 2025

arXiv:2505.20117v1 [math.NT] 26 May 2025

Abstract

We study the 8-rank of class groups of hyperelliptic function fields and show that such 8-ranks are governed by splitting conditions in so-called governing fields. A similar result was proven for quadratic number fields by Stevenhagen, who used a theory of Rédei symbols and Rédei reciprocity to do so. We introduce a version of the Rédei reciprocity law for function fields and use this to show existence of governing fields.

1 Introduction

Since Gauss introduced his genus theory [Gau01], the 2-part of (narrow) class groups of quadratic number fields has been extensively studied. By the Cohen-Lenstra heuristics [CLJ84] we expect only this 2-part to have a predictable structure. To understand the structure of the 2-part, Cohn and Lagarias introduced governing fields and proved their existence for the 2- and 4-rank [CL83]. Stevenhagen gave a proof for the existence of governing fields for the 8-rank [Ste89] using class field theory. More recently, a new proof of their existence [Ste22] was given by using the Rédei symbol, which satisfies the Rédei reciprocity law. This trilinear symbol is named after the Hungarian mathematician who was the first to give a definition [Réd39]. On the existence of a governing field of the 16-rank for quadratic number fields there is only a negative answer by Koymans and Milovic [KM21], under the assumption of a short character sum conjecture.

Similarly to their number field equivalent, class groups of function fields over finite fields show the same structural behaviour in their 2-part, as was first discovered by Artin in his thesis [Art24]. Because of heuristics by Friedman and Washington [FW87] that resemble those of Cohen-Lenstra, this structural behaviour is expected to occur only in the 2-part. This motivates the introduction of governing fields for hyperelliptic function fields, which will be done in this article.

Let q be an odd prime power and consider the rational function field $\mathbb{F}_q(x)$ over the finite field with q elements. A hyperelliptic function field is a quadratic extension $K_D = \mathbb{F}_q(x, \sqrt{D})$ determined by a squarefree polynomial $D \in \mathbb{F}_q[x]$. Its ring of integers $\mathcal{O}_D$ is the algebraic closure of $\mathbb{F}_q[x]$ in K_D and we define $C(D)$ (also denoted by C) as the ideal class group of $\mathcal{O}_D$. Even though we will write $C(D)$ as a multiplicative group, we can make an identification to a vector space over $\mathbb{F}_2$. For $j \in \mathbb{Z}_{\geq 1}$ the 2^j-rank of $C(D)$ is defined as the dimension of the $\mathbb{F}_2$-vector space $C^{2^{j-1}}/C^{2^j}$.

$$r_{2^j} := \dim_{\mathbb{F}_2}\left(C^{2^{j-1}}/C^{2^j}\right) = \dim_{\mathbb{F}_2}\left(C[2^j]/C[2^{j-1}]\right). \tag{1.1}$$

Definition 1.1 (Governing fields)**.** Let q be an odd prime power and $D \in \mathbb{F}_q[x]$ squarefree. A Galois extension $\Omega_{2^j}(D)/\mathbb{F}_q(x)$ is called a *governing field for the* 2^j*-rank of* $C(DP)$ if the Frobenius conjugacy class of an unramified prime P in $\mathrm{Gal}(\Omega_{2^j}(D)/\mathbb{F}_q(x))$ completely determines the 2^k-rank for all $k \leq j$.

In order to obtain results about governing fields we will need explicit methods to determine the 2^j-rank of the class group for various j. Even though this text treats the calculations as a purely algebraic problem for function fields, we want to highlight that similar results can also be obtained through a geometrical interpretation. By relating the class group to the Jacobian of a hyperelliptic curve one obtains additional geometric structure to study the problem [Cor99, Cor01].



*Mathematical Institute, Leiden University and QuSoft, Amsterdam, j.a.stokvis@math.leidenuniv.nl