

Limits to black-box amplification in QMA

Scott Aaronson[1], Phillip Harris[2], and Freek Witteveen[3]

[1]University of Texas at Austin. Supported by the Simons Foundation
[2]University of Bonn
[3]QuSoft & CWI, Amsterdam

arXiv:2509.21131v2 [quant-ph] 9 Oct 2025

Abstract

We study the limitations of black-box amplification in the quantum complexity class QMA. Amplification is known to boost any inverse-polynomial gap between completeness and soundness to exponentially small error, and a recent result (Jeffery and Witteveen, 2025) shows that completeness can in fact be amplified to be doubly exponentially close to 1. We prove that this is optimal for black-box procedures: we provide a quantum oracle relative to which no QMA verification procedure using polynomial resources can achieve completeness closer to 1 than doubly exponential, or a soundness which is super-exponentially small. This is proven by making the oracle separation from (Aaronson, 2009) between QMA and QMA_1 quantitative, using techniques from complex approximation theory.

1 Introduction

The complexity class QMA (Quantum Merlin–Arthur) is a quantum analogue of NP, where given a problem instance, a prover Merlin sends a quantum witness to a polynomial-time quantum verifier Arthur, in order to convince Arthur whether the problem is a yes-instance or a no-instance. Arthur then performs a polynomial-size quantum computation on the witness, after which he accepts or rejects the witness. Quantum complexity classes typically allow some probability of error. There are two parameters quantifying the allowed error: the *completeness* c, the probability with which Arthur accepts a valid witness in the yes-case, and the *soundness* s, the maximum acceptance probability in the no-case. It is well known that the precise choice of these parameters does not matter as long as there is a non-negligible gap: by amplification techniques [KSV02, MW05], one can boost a polynomially small gap $c - s = 1/\operatorname{poly}$ to completeness $c = 1 - 2^{-\operatorname{poly}}$ and soundness $s = 2^{-\operatorname{poly}}$ exponentially close to 1 and 0 respectively. This motivates the canonical definition, which takes $c = \frac{2}{3}$ and $s = \frac{1}{3}$.

A long-standing open question is whether the completeness can in fact be made *perfect*. We denote by QMA_1 the variant of QMA where we take $c = 1$ (and $s = \frac{1}{3}$, or any other constant). The question whether QMA equals QMA_1 is then: can every QMA protocol be modified so that Arthur always accepts a valid witness in the yes-case, while still rejecting no-instances with bounded probability? There are a number of closely related complexity classes which allow for perfect completeness. This is the case for the variant with only classical randomness MA, the variant with a classical proof and quantum verifier QCMA [JKNN12], the variant with multiple rounds QIP [MW05], and the variant with an exponentially small soundness-completeness gap PreciseQMA [FL18].

For QMA, this question has resisted resolution, and one explanation for this is given by Aaronson [Aar09], who gave a barrier to proving $\mathsf{QMA} = \mathsf{QMA}_1$ using black-box techniques. Here 'black-box' refers to an amplification procedure that does not use any properties of the verification circuit, other than those that define it. Concretely, [Aar09] proved that there exists a quantum oracle relative to which $\mathsf{QMA} \neq \mathsf{QMA}_1$. Any proof showing $\mathsf{QMA} = \mathsf{QMA}_1$ would have to break down in the presence of a quantum oracle, i.e. it would have to be quantumly nonrelativizing.

