



Sequential quantum processes with group symmetries

Dmitry Grinko,[1, 2, 3, *] Satoshi Yoshida,[4, *] Mio Murao,[4, 5] and Maris Ozols[1, 2, 3]

[1] *QuSoft, Amsterdam, The Netherlands*
[2] *Institute for Logic, Language and Computation, University of Amsterdam, The Netherlands*
[3] *Korteweg-de Vries Institute for Mathematics, University of Amsterdam, The Netherlands*
[4] *Department of Physics, Graduate School of Science, The University of Tokyo, Hongo 7-3-1, Bunkyo-ku, Tokyo 113-0033, Japan*
[5] *Trans-scale Quantum Science Institute, The University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan*

(Dated: October 9, 2025)

Symmetry plays a crucial role in the design and analysis of quantum protocols. This result shows a canonical circuit decomposition of a $(G \times H)$-invariant quantum comb for compact groups G and H using the corresponding Clebsch–Gordan transforms, which naturally extends to the G-covariant quantum comb. By using this circuit decomposition, we propose a parametrized quantum comb with group symmetry, and derive the optimal quantum comb which transforms an unknown unitary operation $U \in \mathrm{SU}(d)$ to its inverse $U^\dagger$ or transpose U^T. From numerics, we find a deterministic and exact unitary transposition protocol for $d = 3$ with 7 queries to U. This protocol improves upon the protocol shown in the previous work, which requires 13 queries to U.

arXiv:2510.07100v1 [quant-ph] 8 Oct 2025

Symmetry plays an essential role in analyzing quantum systems and quantum information protocols [1–5]. Quantum cloning is one of the most studied protocols in this context, as it highlights the trade-offs between accuracy and resource consumption in quantum state manipulation [6–12]. For $m \to n$ cloners, since the quantum state $U^{\otimes m}|\psi\rangle^{\otimes m}$ should be ideally cloned to a quantum state $U^{\otimes n}|\psi\rangle^{\otimes n}$ for any $U \in \mathrm{U}(d)$ (which is, of course, not possible exactly for $m < n$) and input and output states are invariant under the permutation, the optimal cloning channel satisfies the unitary group covariance and the permutation invariance. In this way, its optimal performance and corresponding protocol are determined. Beyond quantum cloning, symmetry is used in the analysis of various scenarios such as open quantum systems [13, 14], quantum Shannon theory [15–22], quantum error correction [23, 24], and quantum machine learning [25]. References [26, 27] provide a systematic way to utilize the unitary group symmetry to analyze the semidefinite programming, which often appears as the optimization problem in quantum information processing [28, 29]. The quantum Schur transforms and mixed Schur transforms [3, 27, 30–33] provide an efficient implementation of several protocols with the unitary group symmetry (e.g., Refs. [34–36]).

Recently, the quantum supermaps have been developed to analyze the transformation of quantum channels, such as dynamical resource theory [37] and higher-order quantum transformation [38]. The quantum supermaps are implemented by the quantum combs in the quantum circuit, which takes the input state on stream and outputs states accordingly [39]. This circuit structure also describes non-Markovian dynamics [40], and it is also related to the causal structure [41]. The quantum supermaps can be represented as a Choi matrix, and their optimization is often formulated as semidefinite programming [28]. Thus, unitary group symmetry is helpful in analyzing its optimal performance [26]. However, its efficient description in a quantum circuit is not known.

This work shows a canonical circuit decomposition of the $\mathrm{U}(d) \times \mathrm{U}(d)$-invariant quantum comb. This utilizes the (dual) Clebsch–Gordan transforms [3, 27, 30–33] combined with an arbitrary isometry operation acting on the multiplicity space. This construction is extended to $(G \times H)$-invariant quantum comb for compact groups G and H by using the generalized Clebsch–Gordan transforms, which is further extended to the G-covariant quantum comb. By combining this circuit structure with the parametrized quantum comb proposed in Ref. [42], we propose a parameterized quantum comb with group symmetry, which significantly reduces the number of variables to be optimized. This parameterized quantum comb is used to derive the optimal quantum comb for unitary inversion and transposition, which are the tasks to transform n copies of an unknown unitary operation $U \in \mathrm{SU}(d)$ into $U^\dagger$ and U^T, respectively [28, 42–52]. We numerically show that qutrit-unitary transposition is possible with 7 queries of the input unitary U. This is improved over Ref. [52] showing qutrit-unitary transposition with 13 queries.

Quantum comb.— A quantum comb is a quantum circuit composed of n quantum channels $\Psi_i : \mathrm{End}(\mathcal{I}_i \otimes \mathcal{A}_{i-1}) \to \mathrm{End}(\mathcal{O}_i \otimes \mathcal{A}_i)$ for $i \in [n]$ with open slots in between [39], where $\mathcal{I}_i$, $\mathcal{O}_i$, and $\mathcal{A}_i$ are the input, output, and auxiliary systems of the channel Ψ_i, respectively, $\mathcal{A}_0 \simeq \mathcal{A}_n \simeq \mathbb{C}$ are trivial systems, $\mathrm{End}(\mathcal{X})$ represents the set of linear operators on a Hilbert space $\mathcal{X}$, and $[n] := \{1, \ldots, n\}$. It defines a quantum supermap that transforms $n-1$ quantum channels $\Phi_i : \mathrm{End}(\mathcal{O}_i) \to \mathrm{End}(\mathcal{I}_{i+1})$ for $i \in [n-1]$ into a quantum

* These authors contributed equally to this work.
Dmitry Grinko: grinko.dimitry@gmail.com
Satoshi Yoshida: satoshiyoshida.phys@gmail.com