



Optimal Displacement Sensing with Spin-Dependent Squeezed States

Liam J. Bond,[1,2,*] Christophe H. Valahu,[3,4] Athreya Shankar,[5,6] Ting Rei Tan,[3,4] and Arghavan Safavi-Naini[1,2]

[1] *Institute of Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands*
[2] *QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*
[3] *School of Physics, University of Sydney, NSW 2006, Australia*
[4] *Sydney Nano Institute, University of Sydney, NSW 2006, Australia*
[5] *Department of Physics, Indian Institute of Technology Madras, Chennai 600036, India*
[6] *Center for Quantum Information, Communication and Computing, Indian Institute of Technology Madras, Chennai 600036, India*

(Dated: October 31, 2025)

Displacement sensing is a fundamental task in metrology. However, the development of quantum-enhanced sensors that fully utilize the available degrees of freedom in many-body quantum systems remains an outstanding challenge. We propose novel many-body displacement sensing schemes that use spin-dependent squeezed (SDS) states – hybrid spin-boson states whose bosonic squeezed quadrature is conditioned on an auxiliary spin. We prove that SDS states are *optimal*, i.e. their quantum Cramér-Rao bound saturates the Heisenberg limit. We propose explicit measurement sequences that can be readily implemented in systems such as trapped ions. We also introduce a scalable state-preparation protocol and numerically demonstrate the preparation of 8.7 dB of spin-dependent squeezing 15 times faster than the standard approach using second-order sidebands in trapped ions. The potential applications of our sensing protocols range from measuring single-photon scattering to searches for dark matter.

arXiv:2510.25870v1 [quant-ph] 29 Oct 2025

I. INTRODUCTION

Measuring the response of a system to estimate an unknown parameter underpins the development of science and technology. For example, the detection of gravitational waves at the Laser Interferometer Gravitational-Wave Observatory (LIGO) confirmed one of the most important predictions of the general theory of relativity [1, 2], while precision spectroscopy in atomic clocks has revolutionized timekeeping with fractional uncertainties now below 10^{-18} [3, 4]. At the same time, significant efforts have been devoted to using quantum systems for sensing theorized couplings between Standard Model fields and dark matter candidates such as dark photons, millicharged dark matter or axions [5–7]. Such couplings would induce ultra-small bosonic *displacements* of a microwave field [7–11] or a mechanical oscillator [6, 12–16], which could be measured by a sufficiently sensitive displacement sensor.

The sensitivity of the quantum sensor crucially depends on its initial reference state. Using a classical, uncorrelated reference state, the estimation uncertainty is lower bounded by the so-called standard quantum limit (SQL). For displacement sensing, the SQL is defined by bosonic coherent states and is constant, $\mathrm{SQL} \propto \mathcal{O}(1)$. As such, the sensitivity cannot be improved by simply increasing the amplitude of the coherent state. By instead using non-classical reference states and quantum measurement protocols, the estimation uncertainty is ultimately limited by Heisenberg scaling, $\mathrm{HS} \propto 1/\langle \hat{\mathrm{n}} \rangle$, where $\langle \hat{n} \rangle$ is the reference state's average bosonic occupation [17]. This is know as *quantum-enhanced sensing*.

Previous works have considered a range of different bosonic reference states for quantum-enhanced displacement sensing. Examples of states that can achieve sub-SQL sensitivities are Fock states [18, 19] and their (optimal) superpositions [20], phase-twirled squeezed states [21] and two- and four-component cat states [7, 22]. Simultaneous sub-SQL sensing of a displacement's real and imaginary phase-space component can also be achieved using, for example, single-mode grid states [23–25] and two-mode squeezed states [26–28].

However, the aforementioned reference states do not take full advantage of the available degrees of freedom in *many-body* experimental platforms, which feature exquisite control over both spin and bosonic degrees of freedom [29–33]. Such many-body systems are ideal quantum sensors as they offer the possibility of *collective enhancement* for faster reference state preparation, or for amplification of the displacement signal. For example, spin-boson cat states are prepared $\sqrt{N}$ times faster in a trapped-ion system with N ions, while electric field sensing benefits from a $\sqrt{N}$ amplification of the displacement signal when using N ions [15]. While there are existing protocols for single-parameter displacement sensing in many-body systems [15, 34, 35], there is a lack of sensing protocols that utilize the advantages of many-body quantum systems for estimating a displacement's amplitude, or jointly estimating its real and imaginary components.

Here, we solve this problem by introducing many-body displacement sensing schemes using spin-dependent bosonic squeezed states – hybrid spin-boson states whose bosonic squeezed quadrature is conditioned on the state of an auxiliary spin. We prove that such states are *optimal* reference states for both amplitude and joint-parameter displacement sensing, with Quantum Cramér-

* L.J.Bond@uva.nl