

arXiv:2511.13822v1 [quant-ph] 17 Nov 2025

Detection of many-body entanglement partitions in a quantum computer

Albert Rico[1], Dmitry Grinko[2,3,4], Robin Krebs[5], Lin Htoo Zaw[6]
[1] *GIQ - Quantum Information Group, Department of Physics, Autonomous University of Barcelona, Spain*
[2] *QuSoft, Amsterdam, The Netherlands*
[3] *Institute for Logic, Language and Computation, University of Amsterdam, The Netherlands*
[4] *Korteweg-de Vries Institute for Mathematics, University of Amsterdam, The Netherlands*
[5] *Quantum Computing, Technische Universität Darmstadt and*
[6] *Centre for Quantum Technologies, National University of Singapore, 3 Science Drive 2, Singapore 117543*
(Dated: November 19, 2025)

We present a method to detect entanglement partitions of multipartite quantum systems, by exploiting their inherent symmetries. Structures like genuinely multipartite entanglement, m-separability and entanglement depth are detected as very special cases. This formulation enables us to characterize all the entanglement partitions of all three- and four- partite states and witnesses with unitary and permutation symmetry. In particular, we find and parametrize a complete set of bound entangled states therein. For larger systems, we provide a large family of analytical witnesses detecting many-body states of arbitrary size where none of the parties is separable from the rest. This method relies on weak Schur sampling with projective measurements, and thus can be implemented in a quantum computer. Beyond Physics, our results extend to the mathematical literature: we establish new inequalities between matrix immanants, and characterize the set of such inequalities for matrices of size three and four.

I. INTRODUCTION

Over the past decades, a large improvement has been done in the control over quantum systems of increasing size [1, 2]. This advances the development of quantum devices, for which a key feature is the presence of quantum entanglement [3]. Yet, multiple systems can be entangled in different structures, and each is beneficial for different applications. The most resourceful case is arguably genuinely multipartite entanglement, where no subsystem is separable to the rest, and a variety of methods have been derived to detect it [4–6]. However, detecting finer structures is a notorius challenge [7–12].

At the same time, rapid progress in quantum control is increasingly approaching the physical realization of quantum computers [13–15]. Beyond enabling efficient algorithms for classically difficult problems [16–19], quantum computers are also powerful tools for analysing quantum systems, for instance in many-body simulation [20, 21] and quantum chemistry [22, 23]. In particular, quantum hardware is a natural platform where to certify the correlations of quantum systems. While well-established techniques such as randomized measurements are effective for bipartite systems [24–27], less is known for more involved entanglement structures.

Here we develop a technique to detect the entanglement structure of a multipartite system, which is tractable to implement in a quantum computer. By exploiting entanglement symmetries under permutations and unitaries (Observation 2), we detect the entanglement partition a of quantum system (Definition 1). We characterize the entanglement partitions of three- and four-partite invariant systems (Proposition 3 and Table I), and identify the families of bound entangled states therein. Exploiting symmetries further, we derive analytical criteria to certify that no local parties are separable from the rest in many-body systems (Theorem 4). Different methods for practical estimation of the criteria introduced are compared in Table II. Besides contributing to Physics, the results presented lead to new symmetric matrix inequalities, a long-standing problem in the mathematics community (Proposition 5). In particular we characterize the set of immanant inequalities for positive semidefinite matrices of size three and four.




FIG. 1. **Structures of three-partite separability.** Three-partite states can be fully separable (SEP), with separability partition $\kappa = [1, 1, 1]$; biseparable ($\varrho_{AB|C}$, $\varrho_{A|BC}$, $\varrho_{AC|B}$ and their convex combination), with partition $\kappa = [2, 1]$; or genuinely multipartite entangled (GME), with $\kappa = [3]$. We shift symmetric witnesses W detecting biseparable states (green area) by a parameter α_κ, to construct new witnesses detecting states outside of certain separability partitions (blue area).