


arXiv:2005.00607v2 [quant-ph] 13 Sep 2021

Kink dynamics and quantum simulation of supersymmetric lattice Hamiltonians

Jiří Minář,[1,2] Bart van Voorden,[1] and Kareljan Schoutens[1,2]

[1] *Institute for Theoretical Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands*

[2] *QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*

(Dated: September 14, 2021)

We propose a quantum simulation of a supersymmetric lattice model using atoms trapped in a 1D configuration and interacting through a Rydberg dressed potential. The elementary excitations in the model are kinks or (in a sector with one extra particle) their superpartners - the skinks. The two are connected by supersymmetry and display identical quantum dynamics. We provide an analytical description of the kink/skink quench dynamics and propose a protocol to prepare and detect these excitations in the quantum simulator. We make a detailed analysis, based on numerical simulation, of the Rydberg atom simulator and show that it accurately tracks the dynamics of the supersymmetric model.

Introduction. Models of strongly interacting fermions are key to our understanding of condensed matter systems. At the same time, they are notoriously hard to track, even with sophisticated tools ranging from numerical approaches such as quantum Monte-Carlo [1–3] and tensor networks [4, 5] to application of gauge-gravity duality [6]. One strategy to make progress is to consider models with special symmetries. A non-standard but intriguing choice is to consider *supersymmetry* as an explicit symmetry on the lattice [7–12] or as an emergent symmetry [13–15].

$\mathcal{N} = 2$ supersymmetry in a lattice model or a quantum field theory comes with a number of tools, such as the Witten index [16], which facilitate the analysis. Exploiting these tools unveils remarkable features such as extensive ground state degeneracies, a phenomenon dubbed superfrustration [17, 18] which can lead to multicriticality [19].

Despite the supersymmetry, many hard questions remain, such as the nature of the quantum phases in higher spatial dimensions. Here a quantum simulator might provide ingenious insights to these questions. We make a step in this direction and propose such a simulator using arrays of neutral atoms trapped in optical potentials and dressed to their Rydberg state. This is motivated by the high versatility of these platforms [20–29] and by the fact that an off-resonant dressing [30–33] naturally implements the constrained dynamics inherent to the supersymmetric lattice model. Specifically, we consider a so-called M_1 model for spin-less fermions on a 1D chain [8]. As a function of a parameter λ, this model interpolates between a trivial ($\lambda = 0$) and a quantum critical ($\lambda = 1$) phase, the latter connecting to superconformal field theory [8, 34]. The value of the Witten index $W = 2$ indicates the existence of two supersymmetric vacua and points at kinks connecting these two vacua as elementary excitations. Furthermore, in a sector with one particle added, the excitations correspond to the *superpartners* of the kinks, which we call the skinks. We propose a protocol for the (s)kink preparation and detection and solve for their dynamics following a quench (we note a recent study of kink-antikink pair dynamics in a spin chain [35]).




FIG. 1. **(a)** An infinite chain with staggering 11λ can accommodate two ground states $|\mathrm{I}\rangle, |\mathrm{II}\rangle$. The lowest energy states for an open chain of length $L = 3l + 1$ and l particles are kinks $|K_j\rangle$. A supercharge Q acting on a kink creates a particle at the kink location. The blue oval represents the triplet. **(b)** Particle densities in the ground state of an open chain, $L = 3l$, $\lambda = 1$ with an apparent $\mathbb{Z}_3$ pattern highlighted by the light green, dark green and black datapoints. **(c)** Particle (filled symbols) and energy (empty symbols) densities of $|K_1\rangle$ for $\lambda = 0, 0.5, 1$ (blue circles, orange squares, green diamonds). **(d)** Scheme of the proposed realization. Atoms in their electronic ground state $|g\rangle$ tunnel in an optical lattice with spacing r_0 at rate J subject to dressing to a Rydberg state $|r\rangle$. Lower graph shows the dressed potential $W(r)$ for the $|84\mathrm{S}\rangle$ state of ^{6}Li with $\Omega = 2\pi \times 10\,\mathrm{MHz}$, $\Omega/\Delta = 1/10$, $C_6 = 645\,\mathrm{GHz} \cdot \mu\mathrm{m}^6$.

We show that it is identical in both cases and accurately reproduced by the quantum simulator which is a direct consequence and a clear-cut sign of the underlying supersymmetry.

The M_1 model. An $\mathcal{N} = 2$ supersymmetric lattice Hamiltonian for spinless fermions can be defined as

$$H_Q = \{Q, Q^\dagger\}, \tag{1}$$