

2021–
2022

Quantum Channel Marginal Problem

Chung-Yun Hsieh,[1,*] Matteo Lostaglio,[1,2,3] and Antonio Acín[1,4]

[1]*ICFO - Institut de Ciencies Fotoniques, The Barcelona Institute of Science and Technology, 08860 Castelldefels, Spain*
[2]*QuTech, Delft University of Technology, P.O. Box 5046, 2600 GA Delft, The Netherlands*
[3]*Korteweg-de Vries Institute for Mathematics and QuSoft, University of Amsterdam, The Netherlands*
[4]*ICREA-Institució Catalana de Recerca i Estudis Avançats, Lluis Companys 23, 08010 Barcelona, Spain*

(Dated: February 1, 2022)

Given a set of local dynamics, are they compatible with a global dynamics? We systematically formulate these questions as *quantum channel marginal problems*. These problems are strongly connected to the generalization of the no-signaling conditions to quantized inputs and outputs and can be understood as a general toolkit to study notions of quantum incompatibility. In fact, they include as special cases channel broadcasting, channel extendibility, measurement compatibility and state marginal problems. After defining the notion of compatibility between global and local dynamics, we provide a solution to the channel marginal problem that takes the form of a semidefinite program. Using this formulation, we construct channel incompatibility witnesses, discuss their operational interpretation in terms of an advantage for a state-discrimination task, prove a gap between classical and quantum dynamical marginal problems and show that the latter is irreducible to state marginal problems.

arXiv:2102.10926v4 [quant-ph] 31 Jan 2022

Introduction.– A fundamental question in quantum mechanics is whether a given set of local states are compatible with a global one. In other words, can the former be seen as the *marginals* of a global quantum state? This kind of questions are known as *state marginal problems* (SMPs). One of the most prominent examples is the 2-body N-representability problem, where one asks which 2-body reduced density matrices can result as the marginals of a global state of N particles, a problem motivated by the calculation of ground states of 2-body, usually local, Hamiltonians, see for instance [1, 2]. Because of its relevance, the SMP has been studied from many different viewpoints, for instance in the context of entanglement [3, 4] or Bell non-locality detection [5, 6], or by constructing efficient measurement strategies for the estimation of marginal states [7–9]. The SMP also has a classical analogue, where incompatibility appears in the form of frustration due to loops [10].

Whereas the SMP is concerned with static properties encoded in states, the purpose of this work is to understand how compatibility between local and global descriptions extends at the level of dynamics. As we shall see, for the concept of local dynamics to be well-defined, the channels under consideration have to satisfy a generalization of the no-signaling condition to quantum input and outputs considered in Refs. [11–16]. First, however, we introduce the *channel marginal problems* (CMPs) as a natural *dynamical* generalization of the SMP. After this, we provide necessary and sufficient conditions to solve it that can be addressed using semi-definite programming. There exist several previous works that have also considered the CMP [17–22], and we shall see that our framework includes those (and more) within a unified umbrella.

It is convenient for what follows to recall the definition of the SMP, see also Figure 1(a).

Definition 1. (State Marginal Problem) *Consider a global system* S *and a set of local states* $\{\rho_{\mathrm{X}}\}_{\mathrm{X}\in\Lambda}$*, where* Λ *is a collection of subsystems* X *of* S *and each* ρ_{X} *is in* X*. Then a SMP asks whether there exists a global state* ρ_{S} *in* S *compatible with all of them, that is,*

$$\exists\ \rho_{\mathrm{S}} \geq 0 \ \ \textit{such that}\ \ \mathrm{tr}_{\mathrm{S}\setminus\mathrm{X}}(\rho_{\mathrm{S}}) = \rho_{\mathrm{X}},\ \forall\ \mathrm{X}. \tag{1}$$

SMPs always have a trivial solution if the marginals do not overlap, $\rho_{\mathrm{S}} = \bigotimes_{\mathrm{X}\in\Lambda}\rho_{\mathrm{X}}$. The problem becomes much more interesting in the case of overlapping regions. It is again easy to see that the problem has no solution when states ρ_{X} are picked at random. This is because, for the problem to be well-posed, the states ρ_{X} must be compatible in the overlapping regions, that is, their reduced state must be equal. Formally, $\mathrm{tr}_{\mathrm{X}\setminus(\mathrm{X}\cap\mathrm{Y})}(\rho_{\mathrm{X}}) = \mathrm{tr}_{\mathrm{Y}\setminus(\mathrm{X}\cap\mathrm{Y})}(\rho_{\mathrm{Y}})\ \ \forall\ \mathrm{X},\mathrm{Y}\in\Lambda$. These are sometimes called local compatibility conditions, which are necessary and easy to verify but, unfortunately, not sufficient. Still, Eq. (1) is nothing but a set of linear equations on positive operators that can be solved using semi-definite programming (SDP), a standard technique in convex optimization. Note however that the size of the SDP scales exponentially with the number of systems. In fact, the SMP is not expected to have a scalable solution as it is known to be QMA-complete [23].

Channel Marginal Problems.– To formalize the dynamical version of SMPs, first recall that channels correspond to the most general linear operation on quantum states and are represented by completely positive trace-preserving (CPTP) maps [24], denoted by $\mathcal{E}$. One is then tempted to define the CMP as follows: given a set of local channels $\{\mathcal{E}_{\mathrm{X}}\}_{\mathrm{X}\in\Lambda}$, where Λ is a collection of subsystems X of S and each $\mathcal{E}_{\mathrm{X}}$ is a channel in X, the CMP asks whether there exists a global channel $\mathcal{E}_{\mathrm{S}}$ compatible with all of them. The formalization of this compatibility condition, however, is not obvious because the concept of *marginals* for channels is not as straightforward as that for states.

In fact, being an input-output process, the existence of well-defined marginals for a dynamical map requires certain no-signaling conditions. Consider the classical case first. A dynamical map from the the inputs $\mathrm{S}' = \{s'_i\}$ to the outputs $\mathrm{S} = \{s_j\}$ is defined by a stochastic matrix $P_{\mathrm{S}|\mathrm{S}'}$ [25]. Given $\mathrm{X}'\subset\mathrm{S}'$ and $\mathrm{X}\subset\mathrm{S}$, $P_{\mathrm{S}|\mathrm{S}'}$ has a well-defined marginal from X' to X, or simply a reduced map $P_{\mathrm{X}|\mathrm{X}'}$, if and only if for

* chung-yun.hsieh@icfo.eu