


2020–
2024



arXiv:2110.13086v2 [quant-ph] 19 Jul 2022

Quantum Algorithms and Lower Bounds for Linear Regression with Norm Constraints

Yanlin Chen* Ronald de Wolf†

Abstract

Lasso and Ridge are important minimization problems in machine learning and statistics. They are versions of linear regression with squared loss where the vector $\theta \in \mathbb{R}^d$ of coefficients is constrained in either ℓ_1-norm (for Lasso) or in ℓ_2-norm (for Ridge). We study the complexity of quantum algorithms for finding ε-minimizers for these minimization problems. We show that for Lasso we can get a quadratic quantum speedup in terms of d by speeding up the cost-per-iteration of the Frank-Wolfe algorithm, while for Ridge the best quantum algorithms are linear in d, as are the best classical algorithms. As a byproduct of our quantum lower bound for Lasso, we also prove the first classical lower bound for Lasso that is tight up to polylog-factors.

1 Introduction

1.1 Linear regression with norm constraints

One of the simplest, most useful and best-studied problems in machine learning and statistics is *linear regression*. We are given N data points $\{(x_i, y_i)\}_{i=0}^{N-1}$ where $x \in \mathbb{R}^d$ and $y \in \mathbb{R}$, and want to fit a line through these points that has small error. In other words, we want to find a vector $\theta \in \mathbb{R}^d$ of coefficients such that the inner product $\langle \theta, x \rangle = \sum_{j=1}^{d} \theta_j x_j$ is a good predictor for the y-variable. There are different ways to quantify the error ("loss") of such a θ-vector, the most common being the squared error $(\langle \theta, x \rangle - y)^2$, averaged over the N data points (or over an underlying distribution $\mathcal{D}$ that generated the data). If we let X be the $N \times d$ matrix whose N rows are the x-vectors of the data, then we want to find a $\theta \in \mathbb{R}^d$ that minimizes $\|X\theta - y\|_2^2$. This minimization problem has a well-known closed-form solution: $\theta = (X^T X)^+ X^T y$, where the superscript '+' indicates the Moore-Penrose pseudoinverse.

In practice, unconstrained least-squares regression sometimes has problems with overfitting and often yields solutions θ where all entries are non-zero, even when only a few of the d coordinates in the x-vector really matter and one would really hope for a sparse vector θ [SB14, see Chapters 2 and 13]. This situation may be improved by "regularizing" θ via additional constraints. The most common constraints are to require that the ℓ_1-norm or ℓ_2-norm of θ is at most some bound B.[1]

*QuSoft, CWI, the Netherlands. yanlin.chen@cwi.nl

†QuSoft, CWI and University of Amsterdam, the Netherlands. Partially supported by the Dutch Research Council (NWO) through Gravitation-grant Quantum Software Consortium, 024.003.037, and through QuantERA ERA-NET Cofund project QuantAlgo 680-91-034. rdewolf@cwi.nl

[1] For ease of presentation we will set $B = 1$. However, one can also set B differently or even do a binary search over its values, finding a good θ for each of those values and selecting the best one at the end. Instead of putting a hard upper bound B on the norm, one may also include it as a penalty term in the objective function itself, by just