

2019–
2021





2021–
2025



arXiv:2207.10097v3 [quant-ph] 7 Feb 2024

Complexity of the Guided Local Hamiltonian Problem: Improved Parameters and Extension to Excited States

Chris Cade[1,3], Marten Folkertsma[2], and Jordi Weggemans[2]

[1]QuSoft & University of Amsterdam (UvA), Amsterdam, the Netherlands
[2]QuSoft & CWI, Amsterdam, the Netherlands
[3]Fermioniq, Amsterdam, the Netherlands

February 8, 2024

Abstract

Recently it was shown that the so-called *guided local Hamiltonian* problem – estimating the smallest eigenvalue of a k-local Hamiltonian when provided with a description of a quantum state ('guiding state') that is guaranteed to have substantial overlap with the true groundstate – is BQP-complete for $k \geq 6$ when the required precision is inverse polynomial in the system size n, and remains hard even when the overlap of the guiding state with the groundstate is close to a constant $(\frac{1}{2} - \Omega(\frac{1}{poly(n)}))$.

We improve upon this result in three ways: by showing that it remains BQP-complete when i) the Hamiltonian is 2-local, ii) the overlap between the guiding state and target eigenstate is as large as $1 - \Omega(\frac{1}{poly(n)})$, and iii) when one is interested in estimating energies of excited states, rather than just the groundstate. Interestingly, iii) is only made possible by first showing that ii) holds.

1 Introduction

Quantum chemistry is generally regarded as one of the most promising applications of quantum computers [Aar09, BBMC20]. To obtain information about the physical and chemical properties of quantum systems, one usually needs to estimate their spectral properties. For example, in order to understand chemical reaction processes, it is often necessary to know the relative energies of intermediate states along a particular reaction pathway. By comparing these energies, one can deduce which sequence of molecular transformations is most likely to occur in reality. Such energies are usually estimated via some computational method from the *electronic structure Hamiltonian* associated to the system. The accuracy to which these energies are known is tremendously important: typically, in order to confidently distinguish between several reaction mechanisms, one would like to have an accuracy that is smaller than the so-called *chemical accuracy*, which is about 1.6 millihartree.[1] Since in chemistry the norm of the Hamiltonian is allowed to scale polynomially in the number of particles and local dimension per particle, obtaining chemical accuracy corresponds to obtaining inverse polynomial precision when one considers (sub-)normalized Hamiltonians.[2]

The problem of estimating ground- and excited-state energies of the electronic structure Hamiltonian, without any additional information to help us, is known to be QMA-hard [OIWF21].[3] Assuming that QMA $\neq$ BQP, this suggests that estimating energies of physical Hamiltonians is a hard problem even for quantum computers. Hence, to better understand when and how quantum

[1]This quantity, which is $\approx$1 kcal/mol, is chosen to match the accuracy achieved by thermochemical experiments.
[2]By normalized Hamiltonian, we mean a Hamiltonian H such that $\|H\| \leq 1$, where $\|\cdot\|$ denotes the operator norm.
[3]QMA is the set of problems for which a YES-instance can be verified efficiently with a quantum computer, and can be thought of as the quantum analogue of the class NP.