



2019–
2023

# The minimal canonical form of a tensor network

Arturo Acuaviva[1], Visu Makam[2], Harold Nieuwboer[3], David Pérez-García[1], Friedrich Sittner[1], Michael Walter[4], and Freek Witteveen[5]

[1]Departamento de Análisis Matemático y Matemática Aplicada, Universidad Complutense de Madrid, Spain
[2]Radix Trading Europe B.V., Amsterdam, The Netherlands
[3]Korteweg-de Vries Institute for Mathematics and QuSoft, University of Amsterdam, The Netherlands
[4]Faculty of Computer Science, Ruhr University Bochum, Germany
[5]Department of Mathematical Sciences and QMATH, University of Copenhagen, Denmark

arXiv:2209.14358v1 [quant-ph] 28 Sep 2022

**Abstract**

Tensor networks have a gauge degree of freedom on the virtual degrees of freedom that are contracted. A *canonical form* is a choice of fixing this degree of freedom. For matrix product states, choosing a canonical form is a powerful tool, both for theoretical and numerical purposes. On the other hand, for tensor networks in dimension two or greater there is only limited understanding of the gauge symmetry. Here we introduce a new canonical form, the *minimal canonical form*, which applies to projected entangled pair states (PEPS) in any dimension, and prove a corresponding fundamental theorem. Already for matrix product states this gives a new canonical form, while in higher dimensions it is the first rigorous definition of a canonical form valid for any choice of tensor. We show that two tensors have the same minimal canonical forms if and only if they are gauge equivalent up to taking limits; moreover, this is the case if and only if they give the same quantum state for *any* geometry. In particular, this implies that the latter problem is decidable – in contrast to the well-known undecidability for PEPS on grids. We also provide rigorous algorithms for computing minimal canonical forms. To achieve this we draw on geometric invariant theory and recent progress in theoretical computer science in non-commutative group optimization.

## Contents

**1 Introduction** **2**
1.1 Summary of results: a canonical form in any dimension and a fundamental theorem . . . . . 4
1.2 Overview of methods: geometric invariant theory and geodesic convex optimization . . . . . 6
1.3 Organization of the paper . . . . . 6
1.4 Notation . . . . . 6

**2 Preliminaries in geometric invariant theory** **7**

**3 Matrix product states** **9**
3.1 Gauge freedom and canonical forms for uniform MPS . . . . . 9
3.2 The minimal canonical form for uniform MPS . . . . . 11
3.3 Canonical forms for MPS with open boundary conditions . . . . . 16

**4 Projected entangled pair states** **18**
4.1 Definition of uniform PEPS . . . . . 19
4.2 Minimal canonical form . . . . . 21
4.3 Fundamental theorem and invariant theory of uniform PEPS . . . . . 23
4.4 Two-dimensional tensor networks, tilings and topology . . . . . 28
4.5 When does one need the orbit closure? . . . . . 29

**5 Algorithms for computing minimal canonical forms** **33**
5.1 First-order algorithm . . . . . 33
5.2 Relation between approximation errors . . . . . 36
5.3 Second-order algorithm . . . . . 42

**6 Conclusion and outlook** **43**