


2021–
2025

arXiv:2305.01284v4 [quant-ph] 13 Jul 2023

Adiabatic ground state preparation of fermionic many-body systems from a two-body perspective

Dyon van Vreumingen[1, 2] and Kareljan Schoutens[1, 2]

[1] *Institute for Theoretical Physics, Universiteit van Amsterdam, The Netherlands*
[2] *QuSoft, Centrum Wiskunde en Informatica, The Netherlands*

(Dated: July 14, 2023)

A well-known method to prepare ground states of fermionic many-body hamiltonians is adiabatic state preparation, in which an easy to prepare state is time-evolved towards an approximate ground state under a specific time-dependent hamiltonian. However, which path to take in the evolution is often unclear, and a direct linear interpolation, which is the most common method, may not be optimal. In this work, we explore new types of adiabatic paths based on the spectral decomposition of the two-body projection of the residual hamiltonian (the difference between the final and initial hamiltonian). The decomposition defines a set of hamiltonian terms which may be adiabatically interpolated in a piecewise or combined fashion. We demonstrate the usefulness of partially piecewise interpolation through examples involving Fermi-Hubbard models where, due to symmetries, level crossings occur in direct (fully combined) interpolation. We show that this specific deviation from a direct path appropriately breaks the relevant symmetries, thus avoiding level crossings and enabling an adiabatic passage. On the other hand, we show that a fully piecewise scheme, which interpolates every hamiltonian term separately, exhibits a worst-case complexity of $O(L^6/\Delta^3)$ as compared to $O(L^4/\Delta^3)$ for direct interpolation, in terms of the number of one-body modes L and the minimal gap Δ along the path. This suboptimality result suggests that only those terms which break necessary symmetries should be taken into account for piecewise interpolation, while the rest is treated with direct interpolation.

I. INTRODUCTION

Quantum computers are currently regarded as a prime candidate for solving problems in condensed matter physics and chemistry that are untractable for classical computers. In particular, since Feynman's observation of the potential of quantum simulation [1], the pioneering work by Lloyd [2] and the invention of quantum phase estimation [3], interest in the deployment of quantum computers as simulators of highly correlated quantum systems has exploded. A large body of work has been established describing techniques for simulating dynamics of many-body systems on a quantum computer [4–9], and these may be combined with quantum phase estimation in order to estimate eigenenergies [10]. A critical question however, to make these methods useful, is how to prepare the states of interest – be it thermal states or eigenstates of the system under investigation – that serve as input to the algorithms that simulate dynamics or compute energies. Although experimental efforts using heuristics such as variational quantum eigensolvers [11–14] have shown great success in preparing such states for systems of fixed size, much remains unknown with regards to "solving" highly correlated systems in general.

A well-known method for preparing approximate ground states of complex systems is adiabatic state preparation, which uses the adiabatic theorem to carry out quantum computation. While originally formulated as a tool to approximate quantum dynamics on large time scales with respect to the inverse energy gap, the adiabatic theorem was reintroduced to attack combinatorial problems [15] and to study many-body systems such as Fermi-Hubbard models [16, 17] and molecules [18–21].

The idea of adiabatic state preparation (ASP) is to prepare an eigenstate $|\psi^{\mathrm{f}}\rangle$ of a "final" hamiltonian H^{f}, starting with an eigenstate $|\psi^{\mathrm{i}}\rangle$ of an "initial" hamiltonian H^{i} which is straightforward to prepare. Given this initial state, one time-evolves the state according to the time-rescaled Schrödinger equation,

$$i\frac{d}{ds}|\psi(s)\rangle = T\,\mathrm{H}(s)|\psi(s)\rangle, \tag{1}$$

where $s = t/T$ is a dimensionless time, and T is the total (physical) evolution time. (We work in units such that $\hbar = 1$.) The evolution is carried out under a time-dependent hamiltonian $\mathrm{H}(s)$ which equals H^{i} at $s = 0$ and H^{f} at $s = 1$. After an evolution with time s, one obtains a state $|\psi^T(s)\rangle = \mathrm{U}^T(s)|\psi(0)\rangle$, where $|\psi(0)\rangle = |\psi^{\mathrm{i}}\rangle$ and $\mathrm{U}^T(s)$ solves eq. 1. By what is known as the *adiabatic theorem*, the state at the end of this evolution, $|\psi^T(1)\rangle$ will be close to the final eigenstate $|\psi^{\mathrm{f}}\rangle$ if T is sufficiently large. One variant of this adiabatic theorem which precisely indicates what "close" and "sufficiently large" mean in this context, is due to Jansen et al. [22]. The statement is that if $\mathrm{H}(s)$ is a hamiltonian defined on the interval $[0, 1]$ which for every $s \in [0, 1]$ has an instantaneous eigenstate $|\psi(s)\rangle$ whose energy is separated from the rest of the spectrum by $\Delta(s) > 0$, then for any $s \in [0, 1]$, the condition

$$T \geq \frac{1}{\delta}\left(\int_0^s \left[\frac{\|\partial_s^2 \mathrm{H}(\sigma)\|}{\Delta^2(s)} + 7\frac{\|\partial_s \mathrm{H}(\sigma)\|^2}{\Delta^3(s)}\right] d\sigma + B\right) \tag{2}$$

where $\|\cdot\|$ denotes the operator norm and B is a boundary term that may be set to zero if $\dot{\mathrm{H}}(0) = \dot{\mathrm{H}}(1) = 0$, is