

2022–
2024

arXiv:2305.03489v3 [quant-ph] 3 May 2024

No-go theorem for entanglement distillation using catalysis

Ludovico Lami,[1,2,3,*] Bartosz Regula,[4] and Alexander Streltsov[5]

[1] *QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*
[2] *Korteweg–de Vries Institute for Mathematics, University of Amsterdam, Science Park 105-107, 1098 XG Amsterdam, the Netherlands*
[3] *Institute for Theoretical Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands*
[4] *Mathematical Quantum Information RIKEN Hakubi Research Team, RIKEN Cluster for Pioneering Research (CPR) and RIKEN Center for Quantum Computing (RQC), Wako, Saitama 351-0198, Japan*
[5] *Centre for Quantum Optical Technologies, Centre of New Technologies, University of Warsaw, Banacha 2c, 02-097 Warsaw, Poland*

The use of ancillary quantum systems known as catalysts is known to be able to enhance the capabilities of entanglement transformations under local operations and classical communication. However, the limits of these advantages have not been determined, and in particular it is not known if such assistance can overcome the known restrictions on asymptotic transformation rates — notably the existence of bound entangled (undistillable) states. Here we establish a general limitation of entanglement catalysis: we show that catalytic transformations can never allow for the distillation of entanglement from a bound entangled state with positive partial transpose, even if the catalyst may become correlated with the system of interest, and even under permissive choices of free operations. This precludes the possibility that catalysis can make entanglement theory asymptotically reversible. Our methods are based on new asymptotic bounds for the distillable entanglement and entanglement cost assisted by correlated catalysts.

The study of quantum entanglement as a resource has been one of the most fundamental problems in the field of quantum information ever since its inception [1]. To utilize this resource efficiently, it is often required to transform and manipulate entangled quantum systems, which leads to the well-studied question of how quantum states can be converted using only local operations and classical communication (LOCC) [2, 3]. The limits of such conversion capability are represented by asymptotic transformation rates: given many copies of an input quantum state ρ, how many copies of a desired target state can we obtain per each copy of ρ? Such rates are particularly important in the context of purifying noisy quantum states into singlets Φ_2, a task known as entanglement distillation, as well as for the reverse task of using such singlets to synthesize noisy quantum states. This leads to the notions of *distillable entanglement* $E_d(\rho)$ [2], which tells us how many copies of Φ_2 we can extract from a given state ρ, and of *entanglement cost* $E_c(\rho)$ [3], which tells us how many copies of a pure singlet are needed to produce ρ.

A phenomenon that can severely restrict our ability to extract entanglement is known as *bound entanglement* [4]: there exist states from which no entanglement can be distilled, even though their entanglement cost is non-zero. A consequence of this is the irreversibility of entanglement theory — after performing a transformation $\rho \to \omega$, one may not be able to realize the reverse process $\omega \to \rho$ and recover all of the supplied copies of ρ. This contrasts with the asymptotic reversibility of theories such as classical and quantum thermodynamics [4–6]. Although reversibility may still hold in some restricted cases (e.g., for all bipartite pure quantum states [3, 5]), and there are even approaches that may enable reversibility by suitably relaxing the restrictions on the allowed physical transformations [7–10], irreversibility is often a fundamental property of the theory of quantum entanglement that may not be easily evaded [11]. It is then important to understand how, if at all, irreversibility can be overcome.

A promising approach to increase the capabilities of entanglement transformations is the use of so-called *catalysts* [12], that is, ancillary systems that can be employed in the conversion protocol, but must eventually be returned in an unchanged state. Although this phenomenon has been shown to be remarkably powerful in the context of single- and many-copy transformations [12–19], it is unknown whether catalysis can enhance asymptotic conversion rates. This motivates in particular an important question: is the use of catalysis enough to facilitate the reversibility of entanglement theory?

In this paper, we close this question by showing that even very permissive forms of catalytic transformations are insufficient to distill entanglement from bound entangled states. Specifically, we consider the representative class of bound entangled states known as positive partial transpose (PPT) states and we show that the catalytically distillable entanglement of any such state is zero, which is strictly less than its catalytic entanglement cost. The result relies on the establishment of a general upper bound on distillable entanglement under catalytic LOCC operations, namely, the relative entropy of PPT entanglement, which was known to be an upper bound only in conventional, non-catalytic protocols [20, 21]. We show that this limitation persists even if one allows the catalyst to build up correlations with the main system, as well as if one allows sets of operations larger than LOCC, in particular all PPT-preserving transformations. This presents a very general limitation on the power of catalytic transformations of entangled states. We additionally study the applications of resource monotones to constraining asymptotic state conversion with catalytic assistance, obtaining a number of bounds that may be of independent interest.

Preliminaries. — We use $\mathrm{SEP}(A : B)$ to denote the set of states σ_{AB} which are separable across the bipartition $A : B$. The notation $\mathrm{PPT}(A : B)$ will be used to denote the set