




2020–
2024

2020–
2024

MONOGAMY OF HIGHLY SYMMETRIC STATES

RENE ALLERSTORFER[1,2], MATTHIAS CHRISTANDL[3], DMITRY GRINKO[1,4], ION NECHITA[5], MARIS OZOLS[1,4,6,7], DENIS ROCHETTE[8,9], AND PHILIP VERDUYN LUNEL[1,2]

Abstract. We investigate the extent to which two particles can be maximally entangled when they are also similarly entangled with other particles on a complete graph, focusing on Werner, isotropic, and Brauer states. To address this, we formulate and solve optimization problems that draw on concepts from many-body physics, computational complexity, and quantum cryptography. We approach the problem by formalizing it as a semi-definite program (SDP), which we solve analytically using tools from representation theory. Notably, we determine the exact maximum values for the projection onto the maximally entangled state and the antisymmetric Werner state, thereby resolving long-standing open problems in the field of quantum extendibility. Our results are achieved by leveraging SDP duality, the representation theory of symmetric, unitary and orthogonal groups, and the Brauer algebra.

arXiv:2309.16655v2 [quant-ph] 23 Aug 2024

Contents

1. Introduction 2
1.1. Background 2
1.2. Summary of our results 3
2. Preliminaries 3
2.1. Notation 3
2.2. Schur–Weyl dualities for unitary and orthogonal groups 4
2.3. Werner, isotropic and Brauer states 6
2.4. Jucys–Murphy elements 7
2.5. Restrictions of representations of the Brauer algebra to the symmetric group algebra 7
3. General formalisation of the problem 8
3.1. Dual SDP approach 9
3.2. Automorphism group action and edge-transitive graphs 9
4. K_n-Extendibility 10
4.1. Werner states 10
4.2. Isotropic states 10
4.3. Brauer states 16
4.4. Discussion 22
5. Cyclic graph ($n \to \infty$ limit) and Bell state discrimination 24
5.1. Bell state discrimination in the simultaneous communication setting 24
6. Acknowledgements 26
References 26
Appendix A. K_n-Extendibility of Werner states via primal SDP 29
Appendix B. The dual SDPs for K_n-extendibility 30
B.1. Isotropic states 30
B.2. Brauer states when $q = 0$ 32
B.3. Brauer states with fixed p 32
Appendix C. PPT criterion for Brauer states 32

[1]QuSoft, Amsterdam, The Netherlands
[2]CWI, Amsterdam, The Netherlands
[3]Department of Mathematical Sciences, University of Copenhagen, Copenhagen, Denmark
[4]Institute for Logic, Language and Computation, University of Amsterdam, Amsterdam, The Netherlands
[5]Laboratoire de Physique Théorique, Université de Toulouse, CNRS, Toulouse, France
[6]Korteweg-de Vries Institute for Mathematics, University of Amsterdam, The Netherlands
[7]Institute for Theoretical Physics, University of Amsterdam, The Netherlands
[8]Institut de Mathématiques, Université de Toulouse, Toulouse, France
[9]Department of Mathematics and Statistics, University of Ottawa, Ottawa, Canada
E-mail addresses: rene.allerstorfer@cwi.nl, christandl@math.ku.dk, d.grinko@uva.nl, ion.nechita@univ-tlse3.fr, marozols@gmail.com, denis.rochette@uottawa.ca, philip.verduyn.lunel@cwi.nl.