



2021–
2025

Quantum PCPs: on Adaptivity, Multiple Provers and Reductions to Local Hamiltonians

Harry Buhrman[1], Jonas Helsen[2], and Jordi Weggemans[2]

[1]QuSoft, University of Amsterdam & Quantinuum, Partnership House, Carlisle Place, London
[2]QuSoft & CWI, Amsterdam, the Netherlands

arXiv:2403.04841v3 [quant-ph] 8 Jul 2025

We define a general formulation of quantum PCPs, which captures adaptivity and multiple unentangled provers, and give a detailed construction of the quantum reduction to a local Hamiltonian with a constant promise gap. This reduction turns out to be a versatile subroutine to prove properties of quantum PCPs, allowing us to show:

(i) Non-adaptive quantum PCPs can simulate adaptive quantum PCPs when the number of proof queries is constant. In fact, this can even be shown to hold when the non-adaptive quantum PCP picks the proof indices *uniformly* at random from a subset of all possible index combinations, answering an open question by Aharonov, Arad, Landau and Vazirani (STOC '09).

(ii) If the q-local Hamiltonian problem with constant promise gap can be solved in QCMA, then $\mathsf{QPCP}[q] \subseteq \mathsf{QCMA}$ for any $q \in \mathcal{O}(1)$.

(iii) If $\mathsf{QMA}[k]$ has a quantum PCP for any $2 \leq k \leq \mathrm{poly}(n)$, then $\mathsf{QMA}[2] = \mathsf{QMA}$, connecting two of the longest-standing open problems in quantum complexity theory.

Moreover, we also show that there exist (quantum) oracles relative to which certain quantum PCP statements are false. Hence, any attempt to prove the quantum PCP conjecture requires, just as was the case for the classical PCP theorem, (quantumly) non-relativizing techniques.

Jordi Weggemans: jrw@cwi.nl, https://jordiweggemans.github.io