
2020–
2024


2020–
2024







Towards experimental demonstration of quantum position verification using true single photons

Kirsten Kanneworff,[1,*] Mio Poortvliet,[1] Dirk Bouwmeester,[1,2] Rene Allerstorfer,[3,4] Philip Verduyn Lunel,[5,3,4] Florian Speelman,[3,6] Harry Buhrman,[7,3,6] Petr Steindl,[1] and Wolfgang Löffler[1,†]

[1] *Leiden Institute of Physics, Leiden, The Netherlands*

[2] *Department of Physics, University of California, Santa Barbara, California, USA*

[3] *QuSoft, Amsterdam, The Netherlands*

[4] *CWI, Amsterdam, The Netherlands*

[5] *Sorbonne Université, CNRS, LIP6, France*

[6] *University of Amsterdam, The Netherlands*

[7] *Quantinuum, London, United Kingdom*

The geographical position can be a good credential for authentication of a party, this is the basis of position-based cryptography – but classically this cannot be done securely without physical exchange of a private key. However, recently, it has been shown that by combining quantum mechanics with the speed of light limit of special relativity, this might be possible: quantum position verification. Here we demonstrate experimentally a protocol that uses two-photon Hong-Ou-Mandel interference at a beamsplitter, which, in combination with two additional beam splitters and 4 detectors is rendering the protocol resilient to loss. With this we are able to show first results towards an experimental demonstration of quantum position verification.

arXiv:2502.04125v1 [quant-ph] 6 Feb 2025

Since the geographical location is often a good credential of a party in communications, verification thereof could add a useful layer to communication security – this is the case, for instance, with data centers, banks, government buildings, a lab in a quantum network, or even a satellite. Classically, position verification is only possible securely if a shared private key is established which requires physical contact of the parties [1]. In quantum mechanics, mainly thanks to the no-cloning theorem, this can be avoided [2–5]. The general scheme of quantum position verification (QPV) is shown in Fig. 1: Two verifiers V_0 and V_1 share a private communication channel and aim to confirm the location of a third party, the prover P. The verifiers send classical and quantum information, the prover performs a task and returns classical (and possibly quantum) information. The verifiers use this information and the timing and conclude if the prover was at the claimed position or not. This scheme is one-dimensional but can be extended to higher dimensions [6].

However, it quickly was found that attackers with shared entanglement and exploiting quantum teleportation can break quantum position verification protocols, after first attempts [7–9] a general attack was found [10]. This finding has stimulated broad research into the topic [11–25], and it was found that by including classical-information cryptographic tasks, QPV protocols can be made secure for all practical purposes such that attackers require a very large amount of shared entanglement that does only depend on the amount of classical information used in the QPV protocol [26, 27].




Figure 1. Space-time diagram of a one-dimensional QPV protocol showing the prover (P) centered between two verifiers (V_0 and V_1, solid black) where curly (straight) lines indicate quantum (classical) information exchange. Dashed grey lines show a potential form of attack by two adversaries (A_0 and A_1) positioned around the supposed location of the prover that try to mimic the honest prover responses and are restricted to local operations and classical communication (LOCC). Symbols are explained in the text.

In real-world QPV, the quantum information is sent by photons, and two major loopholes emerge from this: First, photons are susceptible to loss during transmission, which opens up a generic attack strategy since the adversaries can claim loss if their measurements have been performed in the wrong basis, for instance. Therefore, fully loss-tolerant protocols are required [23, 28, 29], the first having been developed in Refs. [11, 25]. We will in-

* kanneworff@physics.leidenuniv.nl

† loeffler@physics.leidenuniv.nl