

arXiv:2503.05647v1 [quant-ph] 7 Mar 2025

Phase estimation with partially randomized time evolution

Jakob Günther,[1] Freek Witteveen,[1,2] Alexander Schmidhuber,[3]
Marek Miller,[1] Matthias Christandl,[1] and Aram W. Harrow[3]

[1] *Department of Mathematical Sciences, University of Copenhagen, Universitetsparken 5, 2100 Copenhagen, Denmark*
[2] *QuSoft and CWI, Science Park 123, 1098 XG Amsterdam*
[3] *Center for Theoretical Physics, MIT, Cambridge, MA, 02139, USA*

(Dated: March 10, 2025)

Quantum phase estimation combined with Hamiltonian simulation is the most promising algorithmic framework to computing ground state energies on quantum computers. Its main computational overhead derives from the Hamiltonian simulation subroutine. In this paper we use randomization to speed up product formulas, one of the standard approaches to Hamiltonian simulation. We propose new partially randomized Hamiltonian simulation methods in which some terms are kept deterministically and others are randomly sampled. We perform a detailed resource estimate for single-ancilla phase estimation using partially randomized product formulas for benchmark systems in quantum chemistry and obtain orders-of-magnitude improvements compared to other simulations based on product formulas. When applied to the hydrogen chain, we have numerical evidence that our methods exhibit asymptotic scaling with the system size that is competitive with the best known qubitization approaches.

I. INTRODUCTION

Among the most promising applications of quantum computing is the calculation of ground state energies in quantum many-body physics and quantum chemistry. While in general it is hard for a quantum computer to estimate the ground state energy, quantum phase estimation (QPE) provides an efficient algorithm when a state with significant overlap to the ground state is given. The complexity of the resulting algorithms for chemistry depends largely on the complexity of simulating the Hamiltonian time evolution as a quantum circuit, as QPE requires time simulation of time $\mathcal{O}(\varepsilon^{-1})$ to compute the energy to accuracy ε. The dominant cost of performing QPE is then given by the cost of Hamiltonian evolution and its scaling with time and system size. State-of-the-art resource estimates for realistic examples from chemistry show that the computational cost of performing QPE is still formidable [1–4]. This means that for realizing useful simulations of quantum chemistry, further reductions in the circuit size required to perform Hamiltonian simulation are necessary. This is particularly relevant for devices with a limited number of logical qubits and limited circuit depth. See [5] for a general discussion of QPE for quantum chemistry.

Deterministic product formulas, also known as Trotterization, have the advantage of being conceptually simple and to require little or no ancilla qubits for their implementation as a quantum circuit. In many situations, their performance can be better than what can be expected from rigorous bounds [1, 6] or can even be proven to have good scaling, e.g. in systems with spatially local interactions [7–9]. There also exist powerful post-Trotter Hamiltonian simulation methods [10, 11]; while they have better scaling, methods based on product formulas are still of great interest due to the small number of ancilla qubits required and their good performance for Hamiltonians of interest [1].

This work is motivated by quantum chemical Hamiltonians. For such a Hamiltonian, defined on an active space of N spatial orbitals (corresponding to $2N$ qubits in second quantization), the number of terms in the Hamiltonian scales as $L = \mathcal{O}(N^4)$. Trotter product formulas have a computational cost which scales with L. While this leads to polynomial scaling in N, the resulting circuits rapidly grow impractically large.

An alternative is provided by *randomized product formulas*, in particular qDRIFT [12]. Here, the number of gates no longer depends on L, but rather on λ, the sum of the interaction strengths in the Hamiltonian. If one wants to simulation Hamiltonian evolution for time t, this requires $\mathcal{O}(\lambda^2 t^2)$ gates. So, while this gets rid of the dependence on L, the downside is that the scaling is quadratic in the time (and λ). In particular, since QPE with target accuracy ε requires Hamiltonian simulation for times up to $\mathcal{O}(\varepsilon^{-1})$, using randomized product formulas leads to phase estimation costs scaling with ε^{-2} rather than the optimal Heisenberg-limited scaling of ε^{-1}. Prior work uses an analysis which suggests that combining QPE with qDRIFT leads to an unfavorable scaling of the total cost with the accuracy of the computation [12]. In particular, a naive analysis leads to a scaling of order ε^{-4} or an undesirable dependence on the probability of error [4]. Using a different randomization scheme, however, as well as a different phase estimation scheme, one can restore the scaling to ε^{-2} [13].

A natural generalization of randomized product formulas is to consider *partially randomized* product formulas [14–16], which are effective if there is a relatively small number of dominant terms in the Hamiltonian, and a long tail of small terms (which are nevertheless too large to be ignored altogether). Such behavior is typical for quantum chemistry Hamiltonians; see Section VI and Appendix C for a detailed discussion.