



arXiv:2506.02579v2 [cond-mat.quant-gas] 21 Oct 2025

Confinement-induced resonances for the creation of quasi-one-dimensional ultracold gases of alkali–alkaline-earth dimers

Lorenzo Oghittu,[1,2] Premjith Thekkeppatt,[3,4] Nirav P. Mehta,[5,1,6] Seth T. Rittenhouse,[7,1,6] Klaasjan van Druten,[3,2] Florian Schreck,[3,2] and Arghavan Safavi-Naini[1,2]

[1] *Institute for Theoretical Physics, Institute of Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands*
[2] *QuSoft, Science Park 123, 1098 XG Amsterdam, The Netherlands*
[3] *Van der Waals-Zeeman Institute, Institute of Physics, University of Amsterdam, 1098 XH Amsterdam, Netherlands*
[4] *NNF Quantum Computing Programme, Niels Bohr Institute, University of Copenhagen, Blegdamsvej 17, 2100 Copenhagen, Denmark*
[5] *Department of Physics and Astronomy, Trinity University, San Antonio, Texas 78212, USA*
[6] *ITAMP, Center for Astrophysics, Harvard & Smithsonian, 60 Garden Street, Cambridge, Massachusetts 02138, USA*
[7] *Department of Physics, the United States Naval Academy, Annapolis, Maryland, 21402, USA*

(Dated: October 22, 2025)

We theoretically investigate the role of confinement-induced resonances (CIRs) in low-dimensional ultracold atomic mixtures in the formation of weakly bound dimers. To this end, we examine the scattering properties of a binary atomic mixture confined by a quasi-one-dimensional (quasi-1D) potential. In this regime, the interspecies two-body interaction is modeled as an effective 1D zero-range pseudopotential, with a coupling strength g_{1D} derived as a function of the three-dimensional scattering length a. This framework enables the study of CIRs in harmonically confined systems, with particular attention paid to the case of mismatched transverse trapping frequencies of the two atomic species. Finally, we consider the Bose-Fermi mixture of ^{87}Rb and ^{87}Sr, and identify values of the experimentally accessible parameters for which CIRs can be exploited to create weakly bound molecules.

I. INTRODUCTION

Ultracold gases of polar molecules have emerged as a rapidly advancing field of research [1]. Owing to their long-range, anisotropic dipole-dipole interactions, these systems offer a highly versatile platform for a wide range of applications, including quantum simulation [2], precision measurement [3, 4], and quantum chemistry [5]. However, the creation of an ultracold gas of dipolar molecules poses significant challenges. Experimentally, two methods for creating ultracold gases of ground-state molecules are commonly employed. The first consists of starting with a thermal gas of molecules and cooling it down via laser techniques [6–8]. Yet, the complicated internal, electronic, rotational, and vibrational structure of the molecules can limit their ability to scatter enough photons, often making laser cooling ineffective for many molecular species [9]. The second method involves starting from an ultracold gas of unbound atoms and associating them to weakly bound molecules through adiabatic passage across a resonance. These weakly bound molecules can then be directly associated to their rovibrational ground state via stimulated Raman adiabatic passage (STIRAP) [10–12]. Forming the initial gas of weakly bound molecules usually requires the system to have a readily accessible broad magnetic Feshbach resonance [13] as seen in many bialkali systems, e.g. K-Rb [10]. Because such broad resonances may not be available in some ultracold mixtures, for example due to a weak magnetic response of the atoms or the resonance appearing at very large magnetic fields [14], a different approach becomes necessary.

In this work, we theoretically investigate the use of a different kind of resonance as a pathway for creating weakly bound molecules: the confinement-induced resonance (CIR) [15–21]. CIRs arise in ultracold gases when atoms experience a strong confinement along one or more directions that effectively reduces their dynamics to lower-dimensional regimes. Here, we focus on quasi-one-dimensional (quasi-1D) mixtures of alkali and alkaline-earth atoms. The corresponding heteronuclear dimers constitute a particularly interesting system thanks to their sizable electric and magnetic dipole moments [22]. Unlike bi-alkali dimers, however, the formation of these molecules via magneto-association poses considerable challenges. This is primarily due to the non-magnetic ground state of alkaline-earth atoms, which is responsible for exceedingly narrow Feshbach resonances [23–25]. In this context, the use of CIRs emerges as an appealing alternative. Taking into consideration the relevant Bose-Fermi mixture of ^{87}Rb and ^{87}Sr confined by a quasi-1D harmonic potential (see Fig. 1), we perform a numerical analysis to characterize the emergence of CIRs. We emphasize that the formation of molecules via two-body CIRs requires a coupling between the center of mass and relative motion, as established in Ref. [19]. In the present case, such coupling is introduced by the mismatch in trapping frequencies between the two atomic species, giving rise to narrow inelastic resonances associated with excited center-of-mass bound states. In light of this mechanism, our investigation ultimately aims to identify suitable values of tunable parameters that could support ongoing experimental efforts toward realizing ultracold ^{87}Rb-^{87}Sr dimers.