



Higher spin Richardson-Gaudin model with time-dependent coupling: Exact dynamics

Suvendu Barik,[1,*] Lieuwe Bakker,[1,†] Vladimir Gritsev,[1,2] Jiří Minář,[1,3] and Emil A. Yuzbashyan[4]

[1] *Institute for Theoretical Physics, Universiteit van Amsterdam, Science Park 904, 1098XH Amsterdam, The Netherlands*
[2] *Russian Quantum Center, Skolkovo, Moscow 143025, Russia*
[3] *QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*
[4] *Department of Physics and Astronomy, Center for Materials Theory, Rutgers University, Piscataway, New Jersey 08854 USA*

(Dated: July 16, 2025)

We determine the exact asymptotic many-body wavefunction of a spin-s Richardson-Gaudin model with a coupling inversely proportional to time, for time evolution starting from the ground state at $t = 0^+$ and for arbitrary s. Contrary to common belief, the resulting wavefunction cannot be derived from the spin-1/2 case by merging spins, but instead requires independent treatment for each spin size. The steady state is non-thermal and, in contrast to the spin-1/2 case, does not conform to a natural Generalized Gibbs Ensemble. We show that mean-field theory is exact for any product of a finite number of spin operators on different sites. We discuss how these findings can be probed in cavity QED and trapped ion experiments.

arXiv:2507.10856v1 [quant-ph] 14 Jul 2025

I. INTRODUCTION

Explicit time-dependent Hamiltonians are ubiquitous across all domains of quantum physics. The time dependence may take various forms, such as a quantum quench—a step-like change in parameters [1]—or a continuous, often monotonic variation, as encountered in Landau-Zener (LZ) problems [2, 3], atom-atom and atom-ion collisions [4], and nuclear dynamics [5]. Alternatively, it can be periodic, as in Floquet protocols [6–8], which are used, for example, in studies of time crystals [9–11], the generation of gauge potentials [12], and spin physics [13] in cold atom experiments, as well as in studies of integrable models [14, 15]. The study of such systems remains central to quantum simulation, sensing, and computation, and is closely related to quantum optimal control [16]. Here, the goal is to evolve an easily prepared initial state of the system into a desired target state. Designing efficient protocols for such state preparation is an active area of research [17, 18], with the two standard paradigms being discrete (gate-based)[19–22] and continuous time evolution [23, 24].

In most relevant situations, time-dependent many-body Hamiltonians are not amenable to exact treatment. This necessitates further simplifications, such as mean-field analysis [25] or numerical perturbation theory [26], which often limit these models to weak couplings, small Hilbert spaces, or short-time evolutions. As a result, they remain notoriously difficult to solve analytically. In this context, *exactly solvable models* remain an important recourse, serving as reference points for numerical and approximate methods, and offering clear analytical guidance in the design of quantum state preparation protocols. A well-studied class of such models is periodically driven systems, where, by exploiting their underlying Lie-algebraic structure, exact solutions have been constructed in both closed [14, 15, 27] and open [28–30] settings. More generally, it has been conjectured [31] that a *necessary* condition for the exact solvability of a time-dependent quantum Hamiltonian is its quantum integrability, in the sense of possessing nontrivial parameter-dependent commuting partners. A corresponding *sufficient* condition was later formulated [32] as the existence of a non-Abelian gauge field with zero curvature in the space of system parameters. This framework has unified existing solvable models under the broader umbrella of *integrable time-dependent models* [33–40].

An important class of such solvable models comprises Gaudin magnets and their generalizations, such as the Richardson-Gaudin (RG), inhomogeneous Dicke (also known as Tavis-Cummings), and various central spin Hamiltonians. The celebrated BCS theory maps onto the spin-1/2 RG model when expressed in terms of Anderson pseudospins—bilinear combinations of fermionic creation and annihilation operators [41, 42]. In the time-independent case, this model has been solved using a classical version of the Bethe ansatz [43–46]. Introducing time dependence in the Hamiltonian parameters generally breaks integrability. However, for certain special choices—in particular, when the interaction strength is inversely proportional to time—the model remains integrable [32, 33, 38]. Closely related integrable models include generalized bosonic and fermionic Tavis-Cummings Hamiltonians with linear time dependence [34, 37, 40].

Time-dependent RG models also give rise to a variety of exactly solvable multilevel LZ problems that have attracted considerable attention over the past decades [39, 47–51], with applications ranging from ion-atom collisions [52] and transitions between vibrational modes [53] to Rydberg transitions [54–56] and molecular collisions [57].

The dominant focus so far has been on spin-1/2 RG models. The prevailing assumption is that the physics

* s.k.barik@uva.nl
† l.bakker@friam.nl