




A Robust Strontium Tweezer Apparatus for Quantum Computing

Marijn Venderbosch,[1,2,*] Rik van Herk,[1,2,*] Zhichao Guo,[1,2,*] Jesús del Pozo Mellado,[1,2] Max Festenstein,[1,2] Deon Janse van Rensburg,[1,2] Ivo Knottnerus,[1,2,3,4] Yu Chih Tseng,[3,4] Alexander Urech,[3,4] Robert Spreeuw,[3,4] Florian Schreck,[3,4] Rianne Lous,[1,2,†] Edgar Vredenbregt,[1,2] and Servaas Kokkelmans[1,2]

[1] *Coherence and Quantum Technology group, Department of Applied Physics and Science Education, Eindhoven University of Technology, P.O. Box 513, 5600 MB Eindhoven, The Netherlands*

[2] *Center for Quantum Materials and Technologies Eindhoven (QT/e), P.O. Box 513, 5600 MB Eindhoven, The Netherlands*

[3] *Van der Waals-Zeeman Institute, Institute of Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands*

[4] *QuSoft, Science Park 123, 1098 XG Amsterdam, The Netherlands*

(Dated: January 23, 2026)

Neutral atoms for quantum computing applications show promise in terms of scalability and connectivity. We demonstrate the realization of a versatile apparatus capable of stochastically loading a 5×5 array of optical tweezers with single ^{88}Sr atoms featuring flexible magnetic field control and excellent optical access. A custom-designed oven, spin-flip Zeeman slower, and deflection stage produce a controlled flux of Sr directed to the science chamber. In the science chamber, featuring a vacuum pressure of 3×10^{-11} mbar, the Sr is cooled using two laser cooling stages, resulting in $\sim 3 \times 10^5$ atoms at a temperature of $5(1)\,\mu$K. The optical tweezers feature a $1/e^2$ waist of $0.81(2)\,\mu$m, and loaded atoms can be imaged with a fidelity of ~ 0.997 and a survival probability of $0.99^{+0.01}_{-0.02}$. The atomic array presented here forms the core of a full-stack quantum computing processor targeted for quantum chemistry computational problems.

arXiv:2601.16564v3 [physics.atom-ph] 25 Jun 2026

I. INTRODUCTION

Ultracold atoms trapped in arrays of optical tweezers are a promising platform for quantum computing and quantum simulation [1–3], due to their excellent connectivity [4] and scalability [5, 6]. Recently, strontium (Sr) has emerged as an attractive candidate for such tweezer-based platforms [7, 8], driven by its favorable electronic structure that provides access to narrow and ultra-narrow optical transitions utilized in precision metrology [9, 10]. In Sr, the ultra-narrow clock transition can be used for qubit encoding [7, 11], while the Rydberg-S states are accessible via a single-photon transition, offering high Rabi frequencies for high-fidelity entanglement gates [12, 13].

Unlike alkali atoms, Sr exhibits low vapor pressure, which requires a high temperature oven [14] and multiple stages of laser cooling to be implemented in the experimental apparatus. Additionally, multiple laser systems are needed to address broad, narrow and ultra-narrow atomic transitions. Over the years, several Sr setups have been built, demonstrating e.g. continuous loading of a magneto-optical trap (MOT) [15], single-atom preparation [8, 16, 17], motional ground state cooling [18], qubit manipulation [7, 19] and entanglement [12].

We have designed and realized a compact and versatile Sr apparatus capable of stochastically loading a 5×5 array of optical tweezers with single Sr atoms. For robust operation, we opt for a deflection stage instead of the typically used 2-D MOT. The deflection stage based on transverse optical molasses directs a controlled flux of Sr atoms into the science chamber while removing the direct line-of sight between oven and science chamber. This allows us to maintain low vacuum pressure, which benefits long qubit coherence times, while still maintaining a sufficient loading rate for optical tweezer experiments. The deflection stage has a single retro-reflected laser beam that is forgiving with respect to beam alignment. For modularity, compactness, and stability purposes, the laser systems, referenced to a frequency comb, and the optics needed for intensity and frequency control are rack mounted, and the apparatus can run for multiple days without intervention.

In this work, we demonstrate the successful operation of the apparatus for preparing single Sr atoms in tweezers. Our results showcase the readiness of the setup to serve as a basis for single-qubit gate operations and Rydberg-mediated entanglement. In the near-term this apparatus will provide a neutral atom-based backend for the multi-hardware openly accessible platform Quantum Inspire [20, 21]. We are implementing a full-stack approach and currently operate the Rydberg emulator RySP [22] as digital-twin of our apparatus. The paper is organized as follows. The laser systems involved are discussed in section II, followed by the description of the apparatus in section III. The successful demonstration of the apparatus, describing MOT operation, single atom preparation and imaging can be found in section IV.

II. LASER SYSTEMS

The confinement, control, and detection of our strontium atoms depend on 8 continuous wave (CW) lasers

* These authors contributed equally to this work.

† Author to whom correspondence should be addressed: r.s.lous@tue.nl