

Strassen's support functionals coincide with the quantum functionals

Keiya Sakabe
LMU Munich & MCQST
Munich, Germany
keiya.sakabe@lmu.de

M. Levent Doğan
LMU Munich & MCQST
Munich, Germany
mahmut.dogan@lmu.de

Michael Walter
LMU Munich & MCQST
Munich, Germany
University of Amsterdam & *QuSoft*
Amsterdam, The Netherlands
michael.walter@lmu.de

arXiv:2601.21553v2 [cs.CC] 2 Oct 2026

***Abstract*—Strassen's asymptotic spectrum offers a framework for analyzing the complexity of tensors. It has found applications in diverse areas, from computer science to additive combinatorics and quantum information. A long-standing open problem, dating back to 1991, asks whether Strassen's support functionals are universal spectral points, that is, points in the asymptotic spectrum of tensors. In this paper, we answer this question in the affirmative for tensors over complex numbers by proving that the support functionals coincide with the quantum functionals—universal spectral points that are defined via entropy optimization on entanglement polytopes. We obtain this result as a special case of a general minimax formula for convex optimization on entanglement polytopes (and more general moment polytopes). Our formula can be interpreted as a very general "classical-quantum correspondence" analogous to the well-known relation between the Shannon entropy and the von Neumann entropy. Its proof is based on a recent Fenchel-type duality theorem on Hadamard manifolds due to Hirai. In addition to settling Strassen's question, our results yield a unified and simpler approach to several other tensor invariants, including the asymptotic slice rank.**

In more detail, our main results are as follows:

- **A *classical-quantum correspondence* for convex optimization over entanglement polytopes (and other moment polytopes), which are implicitly-defined polytopes that appear in a variety of contexts from algebraic complexity to quantum information: we show that minimization over the entanglement polytope is equivalent to minimization over the support polytopes if the latter is maximized over all basis choices.**
- **As an immediate corollary, we prove that *Strassen's support functionals coincide with the quantum functionals*. This implies that the former are universal spectral points, resolving a long-standing open problem in algebraic complexity. We also provide a novel characterization of the symmetric quantum functionals in the spirit of Strassen's characterization.**
- **We obtain *new geometric proofs* that the quantum functionals are universal spectral points, and that the symmetric quantum functionals are submultiplicative. These proofs avoid the representation-theoretic machinery used in prior work, suggesting that the classical-quantum correspondence could provide a useful tool for constructing new spectral points.**
- **We derive a novel formula for the *asymptotic slice rank* of a tensor as a minimum, over all basis choices, of the *asymptotic vertex cover number* of associated hypergraphs, which also generalizes to the weighted case.**
- **We obtain direct and uniform new proofs that the G-stable rank and the non-commutative rank of tensors can be expressed as optimizations over the moment polytope, and a general method for obtaining such characterizations for other tensor invariants.**

This research is supported by the European Union (ERC Grant SYMOPTIC, 101040907), by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation, 556164098), and by the Deutsche Forschungsgemeinschaft under Germany's Excellence Strategy - EXC-2111 - 390814868.

I. Introduction

Entropy plays a central role in computer science, mathematics, physics, and many other fields of sciences. Most famous of all is the *Shannon entropy* of a probability distribution $p \in \mathbb{R}^n$:

$$H(p) = -\sum_{\ell=1}^{n} p_\ell \log_2 p_\ell,$$

which measures the "uncertainty" or "information" of the distribution. It originates in statistical physics, going back to the works of Boltzmann and Gibbs, and plays a foundational role in information theory. Moving from classical to quantum information, probability distributions get replaced by quantum states or density matrices ρ, which are positive semidefinite matrices of trace one. While the entries of a density matrix are complex numbers, their eigenvalues form a probability distribution, and hence we can compute their Shannon entropy. The result is called the *von Neumann entropy*:

$$S(\rho) = -\operatorname{tr} \rho \log_2 \rho = H(\operatorname{spec} \rho),$$

where $\operatorname{spec} \rho$ denotes the eigenvalues of the density matrix ρ sorted (for definiteness) in nonincreasing order. It plays a similarly distinguished role in quantum information theory, but also appears in other areas of computer science. Of course, the Shannon and the von Neumann entropy are tightly connected. One well-known but slightly non-obvious fact is the following:

$$S(\rho) = \min_{u \text{ unitary}} H(\operatorname{diag}(u\rho u^\dagger)), \tag{I.1}$$

where $\operatorname{diag}(\cdot)$ sends a matrix to its diagonal entries—in other words, the right-hand side optimizes the Shannon entropy of the diagonal entries, which form a probability distribution, with respect to an arbitrary basis. This basic result has an important interpretation: in quantum information, classical information can be extracted from quantum states by so-called *measurements* in an arbitrary basis, the outcomes of which are given by the diagonal entries in that basis. Thus, Eq. (I.1) states that the classical entropy is always at least as large