



Entropy from Entanglement in Quantum State Reduction

Lisa Lenstra[1,2,*] and Jasper van Wezel[1,†]

[1] *Institute for Theoretical Physics Amsterdam, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands*

[2] *QuSoft, Science Park 123, 1098 XG Amsterdam, The Netherlands*

The Von Neumann entropy of reduced states is a measure of bipartite entanglement. Despite its name, the entanglement entropy cannot by itself be used as a resource for creating thermodynamic heat flows. In order to extract heat from an entangled pure state, it first needs to be converted into a stochastically mixed state by a process of quantum state reduction. Here we show that even in a system with only two degrees of freedom, for which bipartite entanglement is the sole form of entanglement available, the entanglement entropy cannot be converted into thermodynamic entropy in a one-to-one fashion. Moreover, we show that the stochastic dynamics which is necessarily present in any realistic model of quantum state reduction, allows for multiple definitions of entropy. We indicate why quantum state reduction does not allow construction of a perpetuum mobile, despite some measures of entropy evolving non-monotonically during its dynamics. Finally, we relate the different measures of entropy to the information they contain about quantum entanglement and extractable heat, and show that models of quantum state reduction based on physical, correlated stochastic driving forces give rise to observable thermodynamic signatures of quantum state reduction that can be unambiguously distinguished from environment-induced dephasing.

arXiv:2605.00485v1 [quant-ph] 1 May 2026

I. INTRODUCTION

Quantum entanglement and thermodynamic entropy are closely related concepts. In fact, one of the most used measures for bipartite entanglement in quantum systems is known as the (reduced) Von Neumann entropy [1–4]. However, to create a statistical ensemble with non-zero thermodynamic entropy starting from a collection of identical pure entangled quantum states, a physical operation is required. The quantum states need to be measured, in order to project them onto a set of unentangled basis states with Born rule probabilities. Only after such measurement does the ensemble contain thermodynamic entropy that can be used to, for example, create a flow of heat –using the second law– or to do work –using the first law at constant internal energy [5, 6]. The reduction of a pure initial state to a true statistical mixture, rather than tracing a pure state over environmental degrees of freedom to obtain a mixed effective description, can be described by models of objective quantum state reduction [7–10].

Several such models have been proposed as solutions to the measurement problem of quantum mechanics, but to date none have been experimentally verified or falsified, because of the extremely challenging conditions required for probing their predicted dynamics [10–16]. Here, we show that the conversion of quantum entanglement into thermodynamic entropy follows different evolutions for distinct models of quantum state reduction. The generation of thermodynamic entropy thus contains signatures of the way that quantum measurement unfolds, and can be used to distinguish between different types of models for objective quantum state reduction. Moreover, because changes in thermodynamic entropy can be used to generate flows of heat or to do work, the creation of entropy during measurement can be used to constrain models of objective quantum state reduction by requiring that they do not allow any form of perpetuum mobile. We show that this requirement is satisfied for several large classes of models, despite some of them not conserving the average weight of wavefunction components (i.e. their dynamics not being a Martingale process [17, 18]), and some measures for entropy evolving non-monotonically during quantum state reduction.

Finally, we show that multiple, inequivalent ways of defining statistical ensembles are possible for entangled quantum systems undergoing measurement dynamics. We use these to show that regardless of the way in which quantum measurement works, the entanglement entropy in the initial state is never converted into thermodynamic entropy in a one-to-one fashion, indicating that entanglement and entropy are not interchangeable physical properties.

II. QUANTUM STATE REDUCTION

All models for objective quantum state reduction propose generalisations of Schrödinger's equation, whose effect is unobservable at microscopic scales while it dominates the dynamics of macroscopic objects [8–11, 17]. The result on everyday scales is a fast dynamical reduction of generic quantum superpositions to classical states. Importantly, the proposed dynamics affects individual systems, without the need to average over any environmental degrees of freedom, which makes it distinct from environment-induced decoherence [9, 10, 14, 19]. To serve as a solution to the quantum measurement problem, the probability of obtaining any particular classical state in

* e.i.lenstra@uva.nl
† vanwezel@uva.nl