



Quantum Fourier transform for the symmetric group

Carli Bruinsma[1,2], Dmitry Grinko[1,2,3], Maris Ozols[1,2,3]

[1]QuSoft, Amsterdam, The Netherlands
[2]Institute for Logic, Language and Computation, University of Amsterdam, The Netherlands
[3]Korteweg-de Vries Institute for Mathematics, University of Amsterdam, The Netherlands
Email addresses: carlibruinsma@gmail.com, d.grinko@uva.nl, marozols@gmail.com

arXiv:2608.28569v2 [quant-ph] 17 Sep 2026

Abstract

Quantum Fourier transforms (QFT) for general groups were recognized to be fundamental already early in the field. A canonical example of non-abelian QFT for the symmetric group was outlined by Beals (1997). Later, a more detailed analysis of this algorithm was carried out by Kawano and Sekigawa (2016). In this paper, we revisit that construction. After a careful analysis, we revise their gate complexity to $\widetilde{\mathcal{O}}(n^{3.5})$ and circuit depth to $\widetilde{\mathcal{O}}(n^3)$. Moreover, we observe that their construction is not optimal in the choice of transversal elements, so we propose simpler realization of the symmetric group QFT.

Contents

1 Introduction **2**
1.1 Results and their relation to previous work . . . 3

2 Preliminaries and notation **4**
2.1 The symmetric group and its irreducible representations . . . 4
2.2 Chain of subgroups and multiplicity-free restrictions for S_n . . . 4
2.3 Bratteli diagram for S_n . . . 5
2.4 Young–Yamanouchi basis for S_n . . . 5
2.5 Addable and removable cells, Manhattan distances and contents . . . 6
2.6 Yamanouchi words . . . 7

3 Fourier transform along a multiplicity-free subgroup chain **8**
3.1 Regular representations and the Fourier transform . . . 8
3.2 Transversals and induced representations . . . 8
3.3 Induction along the chain of subgroups . . . 10
3.4 Relative path encoding . . . 10

4 Constructing $\mathrm{U}_{\mathrm{Ind}}$ for S_n: Beals approach **12**
4.1 General approach: Looping through transversal elements . . . 12
4.2 Implementation for S_n . . . 13
4.3 Complexity . . . 14

5 Constructing $\mathrm{U}_{\mathrm{Ind}}$ for S_n: Mackey approach **14**
5.1 Mackey theory: Background . . . 15
5.2 Mackey theory and transversals for S_n . . . 16
5.3 The embedding operation . . . 20
5.4 Recovering Kawano–Sekigawa . . . 23
5.5 Implementation of $\mathrm{A}_{\lambda,\widetilde{\lambda}}$ . . . 25
5.6 Complexity . . . 25