



Optimal Ground-State Preparation with a Guiding State

Stacey Jeffery* Rolando D. Somma† Freek Witteveen‡ Ronald de Wolf§

arXiv:2609.38091v2 [quant-ph] 5 Oct 2026

Abstract

Suppose a Hamiltonian H has a unique ground state $|\psi_0\rangle$ with an eigenvalue E_0, and we have an estimate $\tilde{E}_0$ such that $|\tilde{E}_0 - E_0| \leq \delta$, and there is a gap of at least 3δ between E_0 and all other eigenvalues. Suppose we have a unitary A available that can produce a "guiding state" $A|0\rangle$ that has overlap at least γ with $|\psi_0\rangle$. We show how to obtain an η-approximation of $|\psi_0\rangle$ with probability at least $1-\varepsilon$ using $O(\log(1/\varepsilon)/\gamma\delta + \log(1/\eta)/\delta)$ applications of $U = e^{iH}$ its inverse, and $O(\log(1/\varepsilon)/\gamma)$ applications of A and its inverse. We give two different algorithms, one based on interleaving amplitude amplification and error-reduction in the style of [HMdW03], and one using the composition of transducers. This paper is the state-preparation follow-up to our two recent ground-state-energy estimation papers [JW26, SdW26]. Combined with the optimal energy-estimation of [JW26], our results show the claimed upper bound for ground-state preparation. The bound is optimal up to a constant factor in terms of the number of applications of U and $U^\dagger$ if $\varepsilon = \eta$, as follows from [SdW26].

In one of our approaches to ground state preparation, we construct transducers for amplitude amplification and LCU that might be of independent interest.

1 Introduction

We study the problem of preparing the unique ground state $|\psi_0\rangle$ for a given Hamiltonian H. The tools we have available for this are the unitary e^{iH} (which one gets from doing Hamiltonian simulation on H) and a unitary A that prepares a "guiding state" $A|0\rangle$ that is promised to have overlap at least γ with $|\psi_0\rangle$. One may think of $A|0\rangle$ as an "ansatz" for the ground state, based on physical or chemical intuition about the system described by H.

A closely related problem is estimating the ground-state *energy* E_0 associated with the ground state $|\psi_0\rangle$ (i.e., $H|\psi_0\rangle = E_0|\psi_0\rangle$). For the purposes of normalization and to avoid the problem that eigenvalues 0 and 2π are indistinguishable for $U = e^{iH}$, let us assume all eigenvalues are in the interval $[0, \pi]$. This problem of estimating E_0 through a variant of phase estimation is extensively studied in the quantum computing literature. Recently, two of us [JW26] addressed this problem: one can estimate E_0 to within $\pm\delta$ using $O(1/\gamma\delta)$ applications of U and $U^\dagger$. This improves previous work by a factor of $\log(1/\gamma)$, and matches an earlier lower bound of [MdW26] up to a constant factor. At the expense of a factor of $\log(1/\varepsilon)$, one can then reduce the error probability to small ε,

*QLever, QuSoft, CWI and University of Amsterdam, the Netherlands. This work is co-funded by the European Union (ERC, ASC-Q, 101040624).

†Google Quantum AI, Venice, CA 90291, United States.

‡QuSoft and CWI, Amsterdam, the Netherlands

§Google Quantum AI, Venice, CA 90291, United States. Also QuSoft, CWI and University of Amsterdam, the Netherlands. Partially supported by the Dutch Research Council (NWO) through Gravitation-grant Quantum Software Consortium, 024.003.037.