
2024


2017–
2021







arXiv:1809.10751v2 [quant-ph] 10 Dec 2019

Asymptotic performance of port-based teleportation

Matthias Christandl[1], Felix Leditzky*[2,3], Christian Majenz†[4,5], Graeme Smith[2,3,6], Florian Speelman[4,7], and Michael Walter[4,5,8,9]

[1] *QMATH, Department of Mathematical Sciences, University of Copenhagen, Denmark*
[2] *JILA, University of Colorado/NIST, USA*
[3] *Center for Theory of Quantum Matter, University of Colorado Boulder, CO, USA*
[4] *QuSoft, Amsterdam, The Netherlands*
[5] *Institute for Logic, Language and Computation, University of Amsterdam, The Netherlands*
[6] *Department of Physics, University of Colorado Boulder, CO, USA*
[7] *CWI, Amsterdam, The Netherlands*
[8] *Korteweg-de Vries Institute for Mathematics, University of Amsterdam, The Netherlands*
[9] *Institute for Theoretical Physics, University of Amsterdam, The Netherlands*

Abstract

Quantum teleportation is one of the fundamental building blocks of quantum Shannon theory. While ordinary teleportation is simple and efficient, port-based teleportation (PBT) enables applications such as universal programmable quantum processors, instantaneous non-local quantum computation and attacks on position-based quantum cryptography. In this work, we determine the fundamental limit on the performance of PBT: for arbitrary fixed input dimension and a large number N of ports, the error of the optimal protocol is proportional to the inverse square of N. We prove this by deriving an achievability bound, obtained by relating the corresponding optimization problem to the lowest Dirichlet eigenvalue of the Laplacian on the ordered simplex. We also give an improved converse bound of matching order in the number of ports. In addition, we determine the leading-order asymptotics of PBT variants defined in terms of maximally entangled resource states. The proofs of these results rely on connecting recently-derived representation-theoretic formulas to random matrix theory. Along the way, we refine a convergence result for the fluctuations of the Schur-Weyl distribution by Johansson, which might be of independent interest.

Contents

1 Introduction **2**
1.1 Port-based teleportation 2
1.2 Summary of main results 4
1.3 Structure of this paper 8

2 Preliminaries **9**
2.1 Notation and definitions 9
2.2 Representation theory of the symmetric and unitary group 10

*Email: felix.leditzky@jila.colorado.edu
†Email: c.majenz@uva.nl