

2024

2019–
2023




QuSoft is named on page 20 of the paper.

arXiv:2503.22650v1 [math.AG] 28 Mar 2025

Explicit non-free tensors

Maxim van den Berg[1,2], Matthias Christandl[3], Vladimir Lysikov[1], Harold Nieuwboer[3], Michael Walter[1], and Jeroen Zuiddam[2]

[1]Ruhr University Bochum, Bochum, Germany
[2]University of Amsterdam, Amsterdam, Netherlands
[3]University of Copenhagen, Copenhagen, Denmark

Abstract

Free tensors are tensors which, after a change of bases, have free support: any two distinct elements of its support differ in at least two coordinates. They play a distinguished role in the theory of bilinear complexity, in particular in Strassen's duality theory for asymptotic rank. Within the context of quantum information theory, where tensors are interpreted as multiparticle quantum states, freeness corresponds to a type of multiparticle Schmidt decomposition. In particular, if a state is free in a given basis, the reduced density matrices are diagonal. Although generic tensors in $\mathbb{C}^n \otimes \mathbb{C}^n \otimes \mathbb{C}^n$ are non-free for $n \geq 4$ by parameter counting, no explicit non-free tensors were known until now. We solve this hay in a haystack problem by constructing explicit tensors that are non-free for every $n \geq 3$. In particular, this establishes that non-free tensors exist in $\mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^3$, where they are not generic.

To establish non-freeness, we use results from geometric invariant theory and the theory of moment polytopes. In particular, we show that if a tensor T is free, then there is a tensor S in the GL-orbit closure of T, whose support is free and whose moment map image is the minimum-norm point of the moment polytope of T. This implies a reduction for checking non-freeness from arbitrary basis changes of T to unitary basis changes of S. The unitary equivariance of the moment map can then be combined with the fact that tensors with free support have diagonal moment map image, in order to further restrict the set of relevant basis changes.

1 Introduction

Generalizing the notion of a matrix, a k-tensor can be described as a multidimensional array of numbers with k indices $T = (T_{i_1,\dots,i_k})_{i_1,\dots,i_k}$, where the i_ℓ range from 1 to some natural number n_ℓ. They play a pivotal role in algebraic complexity theory (bilinear complexity), quantum information theory (quantum states), combinatorics (cap sets and related problems). In this paper we will restrict our attention to tensors of order 3, as they are of primary relevance for bilinear complexity.

We call any subset $S \subseteq [n]^3$ *free* if any two distinct elements of S differ in at least two coordinates. For any tensor $T \in \mathbb{C}^n \otimes \mathbb{C}^n \otimes \mathbb{C}^n$ let $\operatorname{supp}(T) \subseteq [n]^3$ denote its support, consisting of the multi-indices at which T is non-zero. We call T *free* if there exist $(g_1, g_2, g_3) \in \mathrm{GL}_n(\mathbb{C}) \times \mathrm{GL}_n(\mathbb{C}) \times \mathrm{GL}_n(\mathbb{C})$ such that $\operatorname{supp}((g_1 \otimes g_2 \otimes g_3)T)$ is free; in other words, T has free support after possibly changing the basis on each of the tensor factors. In particular, the tensor is sparse in this basis, in that it has at most n^2 non-zero entries. The class of free tensors play a particularly important role in several parts of the theory of moment polytopes and asymptotic spectra [Fra02, Str05], which we will elaborate on below.

Generic tensors are not free, as can be seen by a dimension argument [CVZ21, Remark 4.19]. In [CGL+21] the dimension of the Zariski-closure of the set of free tensors in $\mathbb{C}^n \otimes \mathbb{C}^n \otimes \mathbb{C}^n$ was determined to be $4n^2 - 3n$. This implies that for every $n \geq 4$ non-free tensors exist in $\mathbb{C}^n \otimes \mathbb{C}^n \otimes \mathbb{C}^n$, but they are dense for $n = 3$. However, no explicit non-free tensors were known until now.