



2024



2019–
2023







QuSoft is named on page 52 of the paper.

Computing moment polytopes — with a focus on tensors, entanglement and matrix multiplication

Maxim van den Berg[1,2], Matthias Christandl[3], Vladimir Lysikov[1], Harold Nieuwboer[3], Michael Walter[1], and Jeroen Zuiddam[2]

[1]Ruhr University Bochum, Bochum, Germany
[2]University of Amsterdam, Amsterdam, Netherlands
[3]University of Copenhagen, Copenhagen, Denmark

arXiv:2510.08336v1 [math.RT] 9 Oct 2025

Abstract

Tensors are fundamental in mathematics, computer science, and physics. Their study through algebraic geometry and representation theory has proved very fruitful in the context of algebraic complexity theory and quantum information. In particular, moment polytopes have been understood to play a key role. In quantum information, moment polytopes (also known as entanglement polytopes) provide a framework for the single-particle quantum marginal problem and offer a geometric characterization of entanglement. In algebraic complexity, they underpin quantum functionals that capture asymptotic tensor relations. More recently, moment polytopes have also become foundational to the emerging field of scaling algorithms in computer science and optimization.

Despite their fundamental role and interest from many angles, much is still unknown about these polytopes, and in particular for tensors beyond $\mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$ and $\mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$ only sporadically have they been computed. We give a new algorithm for computing moment polytopes of tensors (and in fact moment polytopes for the general class of reductive algebraic groups) based on a mathematical description by Franz (J. Lie Theory 2002).

This algorithm enables us to compute moment polytopes of tensors of dimension an order of magnitude larger than previous methods, allowing us to compute with certainty, for the first time, all moment polytopes of tensors in $\mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^3$, and with high probability those in $\mathbb{C}^4 \otimes \mathbb{C}^4 \otimes \mathbb{C}^4$ (which includes the 2×2 matrix multiplication tensor). We discuss how these explicit moment polytopes have led to several new theoretical directions and results.

Contents

1 Introduction **3**
1.1 Tensors, their moment polytopes, and applications 4
1.2 New algorithms for computing moment polytopes 7
1.3 Explicit moment polytopes of $3 \times 3 \times 3$ and $4 \times 4 \times 4$ tensors 8
1.4 Algorithms for moment polytopes for general group actions 9
1.5 Outlook 9
1.6 Overview of the paper 10