

2024



Adiabatic preparation of many-body quantum states: getting the beginning and ending right

Emil T. M. Pedersen,[1] Freek Witteveen,[2] Klaus Mølmer,[3] and Matthias Christandl[1]

[1] *Department of Mathematical Sciences, University of Copenhagen, Universitetsparken 5, 2100 Copenhagen, Denmark*

[2] *Centrum Wiskunde & Informatica and QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*

[3] *Niels Bohr Institute, University of Copenhagen, Jagtvej 155, 2200 Copenhagen, Denmark*

We present numerical calculations, and simulations performed on a Rydberg atom quantum simulator, of the adiabatic evolution of many-body quantum systems around a quantum phase transition. We demonstrate that the end-to-end transfer error, for a given process duration and dissipative losses, can be suppressed by adopting smooth initial and final scheduling functions for the Hamiltonian. We consider a one-dimensional mixed-field Ising model, as well as a chain of Rydberg atoms, and compare numerical calculations and experimental results for the end-to-end transfer error with different schedule functions. We show, in particular, that if the time dependent Hamiltonian is n times differentiable with vanishing 1st to n^{th} order derivatives in the beginning and in the end, the infidelity with respect to the final adiabatic eigenstate scales as $1/T^{n+1}$ when evolving for time T.

arXiv:2512.17780v1 [quant-ph] 19 Dec 2025

The adiabatic theorem [1] states that a system prepared in an eigenstate of a Hamiltonian H and evolved under a slowly varying Hamiltonian $H(t)$ remains in the corresponding instantaneous eigenstate, up to small non-adiabatic corrections. Adiabatic evolution is therefore widely used for preparing eigenstates of complex many-body Hamiltonians. Several strategies are known to suppress non-adiabatic transitions, including locally slowing the evolution near small gaps [2], choosing favorable paths in parameter space [3], and adding counter-diabatic terms that cancel diabatic couplings [4]. The latter yields a 'shortcut to adiabaticity', where the state is subject to a *modified* Hamiltonian, in order to follow the eigenstate of the original target Hamiltonian. These techniques can be combined with variational and optimal-control methods [5–8].

Here we focus on a different aspect of adiabatic evolution that has been known since early work: the fidelity of the *final* state can be much higher than the fidelity of the instantaneous state during the evolution. In the Landau–Zener problem [9–11], a constant coupling V couples two levels with linearly varying energy difference $\Delta E = \alpha t$. The leading-order non-adiabatic error near the avoided crossing scales as α/V, consistent with the Born–Fock estimates. However, at large positive times the loss of population from the adiabatic state is exponentially small, $\propto \exp(-V/\alpha)$ [12, 13]. Similar exponential suppression appears in other analytically solvable two-level problems and can be understood via superadiabatic basis transformations that absorb low-order non-adiabatic corrections [14]. Crucial to this mechanism is the requirement that the adiabatic eigenstates connect smoothly to the initial and final state, i.e., that the time dependent Hamiltonian evolves in a smooth manner throughout the entire process. Additionally, the derivatives need to vanish at the beginning and end of the Hamiltonian path.

In this work, we investigate the precise relation between the smoothness of the boundary conditions and adiabatic state preparation in many-body systems. We combine three components. We prove a rigorous adiabatic theorem showing that n vanishing time derivatives of the Hamiltonian at the initial and final times give an error bound of $O(\varepsilon^{n+1})$ with $\varepsilon = 1/T$, improving previous rigorous estimates of $O(\varepsilon^n)$. We then study this numerically for a one-dimensional mixed-field Ising model, where we construct schedule functions that control the number of vanishing derivatives at the boundaries while leaving the bulk dynamics largely unchanged. Finally, we study this numerically and experimentally on a Rydberg atom chain implemented on the Aquila neutral-atom quantum simulator [15], where we test these schedule constructions under noise and measurement errors. Our main conclusion is that enforcing smoothness at the beginning and end of the protocol reduces end-to-end infidelity, and that this improvement can be obtained with minimal modification of a given schedule.

I *Adiabatic theorem*

We recall the setting of adiabatic Hamiltonian evolution. Consider an n times continuously differentiable family of Hamiltonians $H(\tau)$, where $0 \leq \tau \leq 1$. We consider time evolution under the Hamiltonian $H(t/T)$ for time $t = 0$ to $t = T$ and write

$$\varepsilon = T^{-1}, \qquad \tau = t/T.$$

Let $|\Phi(\tau)\rangle$ be an eigenstate of $H(\tau)$ with eigenvalue $E(\tau)$, separated from the rest of the spectrum by a finite gap (so there are no level crossings and the state and energy are continuous functions of τ). A true state $|\psi(\tau)\rangle$ evolves according to the Schrödinger equation with initial condition $|\psi(0)\rangle = |\Phi(0)\rangle$. For ε small, the adiabatic theorem guarantees that $|\psi(1)\rangle$ has large overlap with $|\Phi(1)\rangle$.

The error depends, amongst other things, on the number of vanishing derivatives of $H(\tau)$ at $\tau = 0$ and $\tau = 1$. If $H(\tau)$ has n vanishing derivatives at both boundaries, previous rigorous results show an error bound of order $O(\varepsilon^n)$ [16–20]. We extend these results as follows.