


LINEAR PROGRAMMING WITH UNITARY-EQUIVARIANT CONSTRAINTS

DMITRY GRINKO[1] AND MARIS OZOLS[1,2]

Abstract. Unitary equivariance is a natural symmetry that occurs in many contexts in physics and mathematics. Optimization problems with such symmetry can often be formulated as semidefinite programs for a d^{p+q}-dimensional matrix variable that commutes with $U^{\otimes p} \otimes \bar{U}^{\otimes q}$, for all $U \in \mathrm{U}(d)$. Solving such problems naively can be prohibitively expensive even if $p + q$ is small but the local dimension d is large. We show that, under additional symmetry assumptions, this problem reduces to a linear program that can be solved in time that does not scale in d, and we provide a general framework to execute this reduction under different types of symmetries. The key ingredient of our method is a compact parametrization of the solution space by linear combinations of walled Brauer algebra diagrams. This parametrization requires the idempotents of a Gelfand–Tsetlin basis, which we obtain by adapting a general method [DLS18] inspired by the Okounkov–Vershik approach. To illustrate potential applications of our framework, we use several examples from quantum information: deciding the principal eigenvalue of a quantum state, quantum majority vote, asymmetric cloning and transformation of a black-box unitary. We also outline a possible route for extending our method to general unitary-equivariant semidefinite programs.

arXiv:2207.05713v2 [quant-ph] 4 Apr 2023

Contents

1. Introduction 2
1.1. Schur–Weyl duality 3
1.2. Applications of Schur–Weyl duality in quantum computing 3
1.3. Group-covariant quantum channels 3
1.4. Group covariance in different contexts 3
1.5. Local unitary equivariance 4
1.6. Optimization subject to local unitary equivariance 5
1.7. Summary of our main result 6
1.8. Intuition 7
1.9. Open problems 7
1.10. Outline 8
2. Preliminaries 8
2.1. Quantum channels 9
2.2. Unitary equivariance 9
2.3. Semidefinite and linear programming 10
2.4. Motivating problem 10
2.5. Optimization under unitary equivariance 11
2.6. Additional symmetries 11

[1]Institute for Logic, Language, and Computation, University of Amsterdam and QuSoft, Amsterdam, The Netherlands
[2]Korteweg-de Vries Institute for Mathematics and Institute for Theoretical Physics, University of Amsterdam, The Netherlands
E-mail addresses: dmitry.grinko@cwi.nl, marozols@gmail.com.
This research was supported by an NWO Vidi grant (Project No. VI.Vidi.192.109).