



Laser cooling for quantum gases

Florian Schreck [1, 2, *] and Klaasjan van Druten [1, *]

[1] *Van der Waals-Zeeman Institute, Institute of Physics, University of Amsterdam, Science Park 904, 1098XH Amsterdam, The Netherlands*

[2] *QuSoft, Science Park 123, 1098 XG Amsterdam, The Netherlands*

arXiv:2209.01026v1 [cond-mat.quant-gas] 2 Sep 2022

Laser cooling exploits the physics of light scattering to cool atomic and molecular gases to close to absolute zero. It is the crucial initial step for essentially all atomic gas experiments in which Bose–Einstein condensation and, more generally, quantum degeneracy is reached. The ongoing development of laser-cooling methods has allowed more elements to be brought to quantum degeneracy, with each additional atomic species offering its own experimental opportunities. Improved methods are opening new avenues, for example, reaching Bose-Einstein condensation purely through laser cooling as well as the realization of continuous Bose-Einstein condensation. Here we review these recent innovations in laser cooling and provide an outlook on methods that may enable new ways of creating quantum gases.

I. INTRODUCTION

Reaching ever lower temperatures has been a frontier of physics for over a century, as the reduction of thermal excitations renders the quantum behaviour of matter more apparent and controllable, enables low-energy quantum phases to arise and reduces decoherence. Laser cooling allows us to cool atomic and molecular gases from room temperature to the microkelvin regime, where particles move with velocities of as little as centimetres per second [1, 2]. Combined with a trapping mechanism, thousands to billions of particles can be accumulated in a small volume. These laser-cooled gas clouds are exploited for precision measurement, enabling clocks that would only go wrong by one second over the lifetime of the Universe [3], atom interferometers that can accurately measure accelerations [4], molecular spectroscopy to search for new physics [5, 6], quantum memories [7], cold electron beams [8] and more [9, 10]. They are also a crucial stepping stone in nearly all degenerate quantum gas experiments (the exception being experiments based on buffer-gas cooling [11, 12]) thereby enabling the exploration of intriguing few- and many-body quantum physics [13, 14]. The importance of this research direction has been recognized by several Nobel Prizes, in particular those for laser cooling and trapping in 1997 [15–17] and for Bose-Einstein condensation of atomic gases in 2001 [18, 19].

II. BASIC LASER-COOLING SCHEME

To reach quantum degeneracy, the most important quantity to consider is the phase-space density (PSD) of an atomic gas. This represents the number of particles per cubic de Broglie wavelength, and has to reach a value of 1 or higher for a Bose-Einstein condensate (BEC) or a degenerate Fermi gas to form. With an experimental setup such as that shown in Fig. 1a) as the baseline of our discussion, a series of cooling and compression techniques are applied to achieve this goal. In brief, a beam of fast atoms effuses under vacuum from an oven and is slowed in a Zeeman slower by the scattering of photons from a counter-propagating laser beam. The slowed atoms are captured in a magneto-optical trap (MOT), where they are further cooled and compressed by light scattering until this cooling method reaches its limits. To go beyond this, the atoms are transferred into a magnetic or optical dipole trap and cooled by evaporation until quantum degeneracy is reached.

Let us now take a more detailed look at these slowing and cooling stages. The atomic beam is slowed by scattering photons from a counter-propagating laser beam via the absorption of the momentum of the photons, leading to a deceleration of $\sim 1\,\mathrm{cm/s}$ in each scattering event. The optically excited atoms emit photons in random directions, thereby recoiling in the opposite direction, which averages out over many scattering events. The Doppler frequency shift of the laser beam reduces while the atoms are slowing down. To keep the atoms in resonance the Zeeman effect can be used by applying a magnetic field that changes along the atoms' trajectory [1, 2]. Such a Zeeman slower, first demonstrated in 1982 [20] creates an atomic beam with a velocity of a few tens of m/s that arrives in a central ultrahigh vacuum (UHV) chamber in which the quantum gas will be created.

Doppler cooling, first proposed in the 1970s [21, 22], is used to cool the atoms further [1, 2]. Three pairs of counter-propagating laser beams of a few cm diameter are sent along three orthogonal axes onto the centre of the UHV chamber. The laser beams are tuned to a frequency below the atomic transition frequency, such that atoms that travel against them with a few $10\,\mathrm{m/s}$ are Doppler shifted onto resonance. Atoms will therefore mostly scat-

* LC4QG@strontiumBEC.com