



Tolerant testing of stabilizer states with a polynomial gap via a generalized uncertainty relation

Zongbo (Bob) Bao[1], Philippe van Dordrecht[2], and Jonas Helsen[1]

[1] *QuSoft and CWI, Amsterdam, The Netherlands*
[2] *QuSoft and University of Amsterdam, Amsterdam, The Netherlands*

Feb 21, 2025

arXiv:2410.21811v2 [quant-ph] 13 Aug 2025

Abstract

We prove a conjecture of Arunachalam & Dutt [AD24] on the existence of a tolerant stabilizer testing algorithm, and achieve an *exponential* improvement in the parameters of the tester. Key to our argument is a generalized uncertainty relation for sets of Pauli operators, based on the Lovász theta function.

1 Introduction

We consider the problem of stabilizer testing: given copies of a state $|\psi\rangle$ with the promise that it is either ϵ_1-close to a stabilizer state (as measured by state fidelity) or ϵ_2-far away, decide which one is the case. This problem has garnered significant interest in recent years following the introduction of the Bell difference sampling algorithm in [GNW21]. There it was proven that Bell difference sampling gives rise to a one-sided tester (i.e. setting $\epsilon_1 = 1$ and $\epsilon_2 < 1$). Their analysis was improved by [GIKL24], who proved that Bell difference sampling gives rise to a tester provided that $\epsilon_2 \leq C\epsilon_1^6$. However, they also required that ϵ_1 was larger than some critical threshold. This latter requirement was relaxed by [AD24], allowing for arbitrary ϵ_1 (this regime is called tolerant testing). However, their argument only obtained a weaker gap relation of the form $\epsilon_2 \leq \exp\{-O(1/\epsilon_1^c)\}$, and furthermore it turned out their argument relied on an unproven statement in additive combinatorics, making their ultimate result conjectural (see Conjecture 1.2 in [AD24][1]).

In this paper we both sidestep their need for a conjecture, and improve their gap relation to a *polynomial* one. This is the polynomial gap mentioned in the title. Concretely we prove the following result:

Theorem 1. *Let $\epsilon_1, \epsilon_2 \in [0,1]$ and let $|\psi\rangle$ be a state with stabilizer fidelity*

$$\mathcal{F}_S(|\psi\rangle) := \max_{|S\rangle \in \mathrm{STAB}_n} |\langle\psi|S\rangle|^2, \tag{1}$$

either (1) larger than ϵ_1 or (2) smaller than ϵ_2. If $\epsilon_2 \leq C'\epsilon_1^{672}$ for some constant C', then $O(\epsilon_1^{-12})$ rounds of Bell difference sampling can distinguish between case (1) and case (2) with probability greater than $2/3$.

[1] More specifically the second version currently on the arXiv. The first version (lemma 4.17) contains contains a misstatement of a result from [Vio11], leading to the requirement of Conjecture 1.2 in the second version.