

Measurement of the g factor of ground-state ^{87}Sr at the parts-per-million level using co-trapped ultracold atoms

Premjith Thekkeppatt,[1,*] Digvijay,[1] Alexander Urech,[1,2] Florian Schreck,[1,2] and Klaasjan van Druten[1,2,†]

[1] *Van der Waals-Zeeman Institute, Institute of Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands*

[2] *QuSoft, Science Park 123, 1098XG Amsterdam, The Netherlands*

(Dated: accepted 8 October 2025; this version compiled: November 4, 2025)

We demonstrate nuclear magnetic resonance of optically trapped ground-state ultracold ^{87}Sr atoms. Using a scheme in which a cloud of ultracold ^{87}Rb is co-trapped nearby, we improve the determination of the nuclear g factor, g_I, of atomic ^{87}Sr by more than two orders of magnitude, reaching accuracy at the parts-per-million level. We achieve similar accuracy in the ratio of relevant g factors between Rb and Sr. This establishes ultracold ^{87}Sr as an excellent linear in-vacuum magnetometer. More generally, our work demonstrates how ultracold neutral atoms can be used for the precise determination of highly relevant atomic and nuclear magnetic properties. These results are important for ongoing efforts toward quantum simulation, quantum computation and optical atomic clocks employing ^{87}Sr, and other neutral alkaline-earth and alkaline-earth-like atoms.

arXiv:2504.11242v2 [physics.atom-ph] 3 Nov 2025

Alkaline-earth and alkaline-earth-like atoms are excellent candidates for quantum simulation of many-body physics [1], quantum computing [2] and atom interferometry [3, 4]. They have been pivotal in realizing state-of-the-art optical atomic clocks [5] and molecular clocks [6]. Many of these applications strongly rely on having a singlet electronic ground state and long-lived triplet metastable states. Among these atoms, fermionic ^{87}Sr is especially suited for optical atomic clocks [7–10], and also stands out because it has the largest nuclear spin quantum number, $I = 9/2$. The tenfold degeneracy arising from this large nuclear spin gives rise to SU(10) spin symmetry. ^{87}Sr is thus an ideal platform to study SU(N) spin Hamiltonians with a great degree of control and isolation—for example implementing Kondo lattice models in an optical lattice [11]. Single ^{87}Sr atoms trapped in optical tweezers hold great potential for realizing arrays of higher-dimensional qubits, i.e. qudits, with up to ten states [12–14]. A high degree of control in state preparation and manipulation of the nuclear spin states are among the requirements for realizing these systems.

To control and manipulate the nuclear spin states in ^{87}Sr coherently, a key quantity is the splitting of the energy levels in the presence of a magnetic field, the Zeeman effect. Alkaline-earth(-like) neutral atoms experience weak splittings and/or shifts in an external magnetic field due to the absence of electronic (orbital and spin) angular momentum in the (singlet) electronic ground state. The splitting is determined by the nuclear g factor, g_I. This factor may be written as $g_I = \mu_I(1 - \sigma_d)/\mu_B I$ [15], where μ_B is the Bohr magneton, μ_I is the magnetic moment of the bare nucleus, and σ_d is the diamagnetic shielding factor due to the surrounding electron cloud. For neutral ^{87}Sr atoms, the value of g_I is known experimentally at the 10^{-4} level [16, 17].

Precision measurement of the g factor of atoms and ions can potentially test quantum electrodynamics (QED) corrections to atomic Hamiltonians involving magnetic interactions. Such measurements are being pursued as a tool for searching for new physics beyond the standard model [18, 19]. By measuring the ratio of the g factor of different isotopes, most of the bound-state QED terms are canceled out, allowing one to probe nuclear effects [20, 21]. The ratio of the microwave clock transitions of ^{87}Rb and ^{133}Cs directly depends on the fine structure constant and the g factor of both atoms and this is used to set limits on the temporal variation of fundamental constants. Improving the accuracy of g-factor values enhances the performance of atomic magnetometers, which rely on the measurement of the Zeeman shift to measure the magnetic field. Atomic magnetometers are often used in a comagnetometer configuration together with noble gases or other isotopes, where the samples overlap spa-




FIG. 1. Schematic of the experimental setup used to measure g_I of ^{87}Sr in the electronic ground state. Clouds of Rb and Sr atoms are confined in respective crossed optical dipole traps, where the (near-)vertical dipole trap beam is common to both traps. The microwave radiation is emitted using a dedicated microwave antenna, whereas the radio frequency radiation is emitted using coils carrying alternating current.