


A random purification channel for arbitrary symmetries with applications to fermions and bosons

Michael Walter* Freek Witteveen†

arXiv:2512.15690v2 [quant-ph] 22 Dec 2025

Abstract

The random purification channel maps n copies of any mixed quantum state ρ to n copies of a random purification of ρ. We generalize this construction to arbitrary symmetries: for any group G of unitaries, we construct a quantum channel that maps states contained in the algebra generated by G to random purifications obtained by twirling over G. In addition to giving a surprisingly concise proof of the original random purification theorem, our result implies the existence of fermionic and bosonic Gaussian purification channels. As applications, we obtain the first tomography protocol for fermionic Gaussian states that scales optimally with the number of modes and the error, as well as an improved property test for this class of states.

1 Introduction

The church of the larger Hilbert space teaches that one can interpret any mixed state as the reduced state of a pure state, and that similarly any process can be purified by the Stinespring extension. Indeed, for any mixed state ρ on a quantum system with Hilbert space $\mathcal{H}$, there exists a pure state $|\psi_\rho\rangle$ on $\mathcal{H} \otimes \mathcal{H}'$, where the reference system $\mathcal{H}'$ is a copy of $\mathcal{H}$, such that the reduced state on $\mathcal{H}$ equals ρ. This purification is not unique: applying an arbitrary unitary $U \in \mathrm{U}(\mathcal{H}')$ to the reference system yields another purification, and in fact any two purifications of ρ differ by such a unitary. This is an extremely useful principle in quantum information theory, where it is often much easier to reason about pure states than about mixed states.

How does one put this creed into practice? Recently, it has been shown that purifications not only exist, but can in fact be constructed in a meaningful way by a *quantum channel* [TWZ25, PSTW25, GML25], see also [SW22, CWZ24]. While no quantum channel can assign a fixed purification to every quantum state, there does exist a *random purification channel* that prepares a purification with a random unitary applied to the reference system. Crucially, it can be extended to multiple copies of the state, such that it does so consistently. That is, the random purification channel $\mathcal{P}_{n,d}$ acts on n copies of a quantum system with Hilbert space $\mathcal{H} = \mathbb{C}^d$, and it has the property that for every state σ on $\mathcal{H}$,

$$\mathcal{P}_{n,d}[\sigma^{\otimes n}] = \int_{\mathrm{U}(d)} \left((I \otimes U)\psi_\sigma^{\mathrm{std}}(I \otimes U^\dagger)\right)^{\otimes n} \mathrm{d}U = \mathop{\mathbf{E}}_{\psi_\sigma \text{ purification of } \sigma} \psi_\sigma^{\otimes n}, \tag{1.1}$$

where $\psi_\sigma^{\mathrm{std}}$ is a fixed (standard) purification of σ.

This construction has useful applications. In particular, it has been applied in [PSTW25] to improve tomography for mixed states. The idea is that if one has a tomography protocol for pure states, then this automatically extends to a tomography protocol for mixed states, as follows:

1. Given n copies of a mixed state σ, apply the random purification channel.
2. Apply pure state tomography on the output of the random purification channel.

*LMU Munich and University of Amsterdam & QuSoft, Amsterdam, michael.walter@lmu.de
†CWI & QuSoft, Amsterdam, f.witteveen@cwi.nl