
2015–
2019




2015–
2019

# The Power Light Cone of the Discrete Bak-Sneppen, Contact and other local processes


Tom Bannink*‡ bannink@cwi.nl

Harry Buhrman*‡ buhrman@cwi.nl

András Gilyén*§ gilyen@cwi.nl

Mario Szegedy¶ mario.szegedy@alibaba-inc.com


April 1, 2019

arXiv:1903.12607v1 [math.PR] 29 Mar 2019


**Abstract**

We consider a class of random processes on graphs that include the discrete Bak-Sneppen (DBS) process and the several versions of the contact process (CP), with a focus on the former. These processes are parametrized by a probability $0 \leq p \leq 1$ that controls a local update rule. Numerical simulations reveal a phase transition when $p$ goes from 0 to 1. Analytically little is known about the phase transition threshold, even for one-dimensional chains. In this article we consider a power-series approach based on representing certain quantities, such as the survival probability or the expected number of steps per site to reach the steady state, as a power-series in $p$. We prove that the coefficients of those power series stabilize as the length $n$ of the chain grows. This is a phenomenon that has been used in the physics community but was not yet proven. We show that for local events $A, B$ of which the support is a distance $d$ apart we have $\mathrm{cor}(A, B) = \mathcal{O}(p^d)$. The stabilization allows for the (exact) computation of coefficients for arbitrary large systems which can then be analyzed using the wide range of existing methods of power series analysis.


## 1 Introduction

In physics, critical behaviour involves systems in which correlations decay as a power law with distance. It is an important topic in many areas of physics and can also be found in stochastic processes on graphs. Often, such systems have a parameter (e.g. temperature) and when it is set to a critical value, the system exhibits critical behaviour. Power series expansion techniques have been used in the physics literature to numerically approximate critical values and associated exponents. It was often observed that the coefficients of such power series stabilize when the system size grows, and we provide a rigorous proof of this for a large class of stochastic processes.

Self-organized criticality is a name common to models where the critical behaviour is present but without the need of tuning a parameter. A simple model for evolution and self-organized criticality was proposed by Bak and Sneppen [BS93] in 1993. In this random process there are $n$ vertices on a cycle each representing a species. Every vertex has a fitness value


*Qusoft, CWI, University of Amsterdam, Science Park 123, 1098 XG Amsterdam, Netherlands.



‡Supported by the NWO Gravitation-grant QSC and NETWORKS-024.002.003, and by EU grant QuantAlgo.

§Supported by ERC Consolidator Grant QPROGRESS and QuantERA project QuantAlgo 680-91-034.



¶Alibaba Quantum Laboratory, Alibaba Group, Bellevue, WA 98004, USA, Also supported by NSF grants 1422102 and 1514164.