

2018–
2020

arXiv:1906.07948v2 [math.CO] 25 Jun 2019

Group-theoretic generalisations of vertex and edge connectivities

Yinan Li [*] Youming Qiao [†]

June 26, 2019

Abstract

Let p be an odd prime. Let P be a finite p-group of class 2 and exponent p, whose commutator quotient $P/[P,P]$ is of order p^n. We define two parameters for P related to central decompositions. The first parameter, $\kappa(P)$, is the smallest integer s for the existence of a subgroup S of P satisfying (1) $S \cap [P,P] = [S,S]$, (2) $|S/[S,S]| = p^{n-s}$, and (3) S admits a non-trivial central decomposition. The second parameter, $\lambda(P)$, is the smallest integer s for the existence of a central subgroup N of order p^s, such that P/N admits a non-trivial central decomposition.

While defined in purely group-theoretic terms, these two parameters generalise respectively the vertex and edge connectivities of graphs: For a simple undirected graph G, through the classical procedures of Baer (Trans. Am. Math. Soc., 1938), Tutte (J. Lond. Math. Soc., 1947) and Lovász (B. Braz. Math. Soc., 1989), there is a p-group P_G of class 2 and exponent p that is naturally associated with G. Our main results show that the vertex connectivity $\kappa(G)$ is equal to $\kappa(P_G)$, and the edge connectivity $\lambda(G)$ is equal to $\lambda(P_G)$. We also discuss the relation between $\kappa(P)$ and $\lambda(P)$ for a general p-group P of class 2 and exponent p, as well as the computational aspects of these parameters.

Keywords: p-groups of class 2, graph connectivity, matrix spaces, bilinear maps

2010 MSC: 20D15, 05C40, 15A69

1 Introduction

The main purpose of this note is to define and explore two natural group-theoretic parameters, which are closely related to vertex and edge connectivities in graphs.

In this introduction, we first introduce the classical procedures of Baer [Bae38], Tutte [Tut47], and Lovász [Lov89] which relate graphs with p-groups of class 2 and exponent p. We then define two group-theoretic parameters. Our main result shows that the vertex and edge connectivities of a graph are equal to the two parameters we defined on the corresponding group respectively. We then compare the two parameters and discuss on their computational aspects.

Since the main goal of this note is to set up a link between graph theory and group theory, we shall include certain background information, despite that it is well-known to researchers in the respective areas.

[*] Centrum Wiskunde & Informatica and QuSoft, Science Park 123, 1098XG Amsterdam, Netherlands (Yinan.Li@cwi.nl). Partially supported by ERC Consolidator Grant 615307-QPROGRESS.

[†] Center for Quantum Software and Information, University of Technology Sydney, Ultimo NSW 2007, Australia. Youming.Qiao@uts.edu.au. Partially supported by the Australian Research Council DECRA DE150100720.