



arXiv:1609.07476v2 [math.CO] 10 Sep 2019

Asymptotic tensor rank of graph tensors: beyond matrix multiplication

Matthias Christandl[1], Péter Vrana[2] and Jeroen Zuiddam[3]

Abstract

We present an upper bound on the exponent of the asymptotic behaviour of the tensor rank of a family of tensors defined by the complete graph on k vertices. For $k \geq 4$, we show that the exponent per edge is at most 0.77, outperforming the best known upper bound on the exponent per edge for matrix multiplication ($k = 3$), which is approximately 0.79. We raise the question whether for some k the exponent per edge can be below 2/3, i.e. can outperform matrix multiplication even if the matrix multiplication exponent equals 2. In order to obtain our results, we generalise to higher order tensors a result by Strassen on the asymptotic subrank of tight tensors and a result by Coppersmith and Winograd on the asymptotic rank of matrix multiplication. Our results have applications in entanglement theory and communication complexity.

1. Introduction

A famous open problem in algebraic complexity theory is the problem of determining the tensor rank of large tensor powers of the 2×2 matrix multiplication tensor. Let $V_1, \ldots, V_k$ be finite-dimensional complex vector spaces and let $\phi \in V_1 \otimes \cdots \otimes V_k$ be a k-tensor. The tensor rank $\mathrm{R}(\phi)$ of ϕ is the smallest number r such that ϕ can be written as a sum of r simple tensors $v_1 \otimes \cdots \otimes v_k \in V_1 \otimes \cdots \otimes V_k$. We define the Nth tensor power $\phi^{\otimes N}$ as the k-tensor obtained by taking the tensor product of N copies of ϕ and grouping such that $\phi \in (V_1^{\otimes N}) \otimes \cdots \otimes (V_k^{\otimes N})$. For any n, let $b_1, \ldots, b_n$ denote the standard basis of $\mathbb{C}^n$. The 2×2 matrix multiplication tensor is the 3-tensor

$$\sum_{i \in \{0,1\}^3} (b_{i_1} \otimes b_{i_2}) \otimes (b_{i_2} \otimes b_{i_3}) \otimes (b_{i_3} \otimes b_{i_1}) \in (\mathbb{C}^2 \otimes \mathbb{C}^2)^{\otimes 3}.$$

[1]QMATH, Department of Mathematical Sciences, University of Copenhagen, Universitetsparken 5, 2100 Copenhagen, Denmark. Email: christandl@math.ku.dk

[2]Department of Geometry, Budapest University of Technology and Economics, Egry József u. 1., 1111 Budapest, Hungary. Email: vranap@math.bme.hu

[3]QuSoft, CWI Amsterdam and University of Amsterdam, Science Park 123, 1098 XG Amsterdam, Netherlands. Email: j.zuiddam@cwi.nl