

2018–
2022

A Sublinear-Time Quantum Algorithm for Approximating Partition Functions

Arjan Cornelissen*† Yassine Hamoudi‡

January 10, 2023

arXiv:2207.08643v2 [quant-ph] 7 Jan 2023

Abstract

We present a novel quantum algorithm for estimating Gibbs partition functions in *sublinear time* with respect to the logarithm of the size of the state space. This is the first speed-up of this type to be obtained over the seminal nearly-linear time algorithm of Štefankovič, Vempala and Vigoda [ŠVV09]. Our result also preserves the quadratic speed-up in precision and spectral gap achieved in previous work by exploiting the properties of quantum Markov chains. As an application, we obtain new polynomial improvements over the best-known algorithms for computing the partition function of the Ising model, counting the number of k-colorings, matchings or independent sets of a graph, and estimating the volume of a convex body.

Our approach relies on developing new variants of the quantum phase and amplitude estimation algorithms that return nearly *unbiased* estimates with *low variance* and *without destroying* their initial quantum state. We extend these subroutines into a nearly unbiased quantum mean estimator that reduces the variance quadratically faster than the classical empirical mean. No such estimator was known to exist prior to our work. These properties, which are of general interest, lead to better convergence guarantees within the paradigm of simulated annealing for computing partition functions.

1 Introduction

The Boltzmann–Gibbs distribution is a paradigmatic tool for modeling systems that obey the principle of maximum entropy. It arises in several fields of research such as statistical mechanics [Geo11; FV17; Sin82], economic modeling [DY00], image processing [GG84; Bes86], statistical learning theory [Cat04], etc. The probability assigned by the Gibbs distribution to each possible configuration of a system is inversely proportional to the exponential of its energy multiplied by the inverse temperature. Mathematically, for a classical Hamiltonian $H : \Omega \to \{0, \dots, n\}$ of degree n specifying the energy level of each configuration x, the Gibbs distribution at inverse temperature β is given by $\pi_\beta(x) = \frac{1}{Z(\beta)} e^{-\beta H(x)}$ where the normalization factor

$$Z(\beta) = \sum_{x \in \Omega} e^{-\beta H(x)} \tag{1}$$

is called the *partition function*. While it is often straightforward to evaluate the partition function at high temperature (when $\beta = 0$, it is just the number of possible configurations), the low-temperature regime captures ground state properties that are challenging to compute. For instance, $Z(\infty)$ can represent the cardinalities of exponentially large combinatorial structures (such as the number of colorings of a graph [Jer95; ŠVV09], the volume of convex bodies [DFK91;

*QuSoft, University of Amsterdam. arjan.cornelissen@cwi.nl
†IBM Quantum, IBM T.J. Watson Research Center. arjan.cornelissen@ibm.com
‡Simons Institute for the Theory of Computing, University of California, Berkeley. ys.hamoudi@gmail.com