



2020–
2025

2025:7

TheoretiCS

Multidimensional Quantum Walks, with Application to k-Distinctness

Received Sep 27, 2023
Revised Aug 28, 2024
Accepted Oct 28, 2024
Published Mar 4, 2025

Key words and phrases
quantum walk, random walk, k-distinctness, element distinctness, welded trees, quantum algorithm

Stacey Jeffery [a]
Sebastian Zur [a]

a CWI & QuSoft, the Netherlands

ABSTRACT. While the quantum query complexity of k-distinctness is known to be $O(n^{\frac{3}{4}-\frac{1}{4}\frac{1}{2^k-1}})$ for any constant $k \geq 4$ [Belovs, FOCS 2012], the best previous upper bound on the time complexity was $\widetilde{O}(n^{1-1/k})$. We give a new upper bound of $\widetilde{O}(n^{\frac{3}{4}-\frac{1}{4}\frac{1}{2^k-1}})$ on the time complexity, matching the query complexity up to polylogarithmic factors. In order to achieve this upper bound, we give a new technique for designing quantum walk search algorithms, which is an extension of the electric network framework. We also show how to solve the welded trees problem in $O(n)$ queries and $O(n^2)$ time using this new technique, showing that the new quantum walk framework can achieve exponential speedups.

1. Introduction

In the problem of *element distinctness*, the input is a list of n integers, and the output is a bit indicating whether the integers are all distinct, or there exists a pair of integers that are the same, called a *collision*. This problem has been studied as a fundamental problem in query complexity, but also for its relationship to other more practical problems, such as sorting, or *collision finding*, which is similar, but one generally assumes there are many collisions and one wants to find one. In the worst case, element distinctness requires $\Theta(n)$ classical queries [2].

The first quantum algorithm to improve on this was a $O(n^{3/4})$ query algorithm [16], which is a variation of an optimal quantum algorithm for collision finding [15], whose main technique is amplitude amplification [14]. The algorithm of [16] could also be implemented time efficiently,

This work is supported by ERC STG grant 101040624-ASC-Q, NWO Klein project number OCENW.Klein.061, and ARO contract no W911NF2010327. SJ is a CIFAR Fellow in the Quantum Information Science Program. A preliminary version of this article appeared at STOC 2023 [27].