

2022–
2025



arXiv:2307.13730v1 [quant-ph] 25 Jul 2023

Fermionic Hamiltonians without trivial low-energy states

Yaroslav Herasymenko[*,1,2,3], Anurag Anshu[4], Barbara M. Terhal[1,2], and Jonas Helsen[3]

[1]QuTech, TU Delft, P.O. Box 5046, 2600 GA Delft, The Netherlands
[2]Delft Institute of Applied Mathematics, TU Delft, 2628 CD Delft, The Netherlands
[3]QuSoft and CWI, Science Park 123, 1098 XG Amsterdam, The Netherlands
[4]School of Engineering and Applied Sciences, Harvard University, 150 Western Ave., Allston, MA 02134, US

July 27, 2023

Abstract

We construct local fermionic Hamiltonians with no low-energy trivial states (NLTS), providing a fermionic counterpart to the NLTS theorem. Distinctly from the qubit case, we define trivial states via finite-depth *fermionic* quantum circuits. We furthermore allow free access to Gaussian fermionic operations, provided they involve at most $O(n)$ ancillary fermions. The desired fermionic Hamiltonian can be constructed using any qubit Hamiltonian which itself has the NLTS property via well-spread distributions over bitstrings, such as the construction in [ABN22]. We define a fermionic analogue of the class quantum PCP and discuss its relation with the qubit version.

1 Introduction

One of the main problems in quantum complexity is establishing the hardness of quantum computational problems such as the estimation of the ground state energy of a many-body Hamiltonian. If high accuracy is required, –scaling inversely polynomial with the problem size n–, this problem is known to be QMA-complete [KSV02]. It has been an open question whether the estimation of the ground state energy *density* of a Hamiltonian with constant error, would also be QMA-hard [AAV13]. Such a result would mirror the classical PCP theorem, which states that there are classical SAT problems for which it is NP-complete to decide whether all clauses can be satisfied or a constant fraction of the clauses can never be satisfied.

Motivated by this quantum PCP problem, it was conjectured in [FH13] that there exist local sparse qubit Hamiltonians such that any state with sufficiently small energy density cannot be prepared by a constant-depth circuit, hence suggesting that low-energy states have some nontrivial complexity. This "no low-energy trivial state" (NLTS) conjecture was recently proven in [ABN22]. They demonstrate that a family of Hamiltonians based on recently established good quantum LDPC codes [LZ22] has the desired property. While the low energy states of this family of Hamiltonians are non-trivial, their ground state can be efficiently prepared by a Clifford circuit. This means that these Hamiltonians fall short of being genuinely 'complex'. Hence it is natural to extend the definition of "non-trivial" beyond constant-depth circuits to climb up the complexity ladder, see [GG22, WFC23, CCNN23, AGK23] . In particular, [CCNN23] have shown that an

[*]Corresponding author: yaroslav@cwi.nl