



2022–
2024

arXiv:2310.02252v1 [quant-ph] 3 Oct 2023

GELFAND–TSETLIN BASIS FOR PARTIALLY TRANSPOSED PERMUTATIONS, WITH APPLICATIONS TO QUANTUM INFORMATION

DMITRY GRINKO[1], ADAM BURCHARDT[1], AND MARIS OZOLS[1,2]

Abstract. We study representation theory of the partially transposed permutation matrix algebra, a matrix representation of the diagrammatic walled Brauer algebra. This algebra plays a prominent role in mixed Schur–Weyl duality that appears in various contexts in quantum information. Our main technical result is an explicit formula for the action of the walled Brauer algebra generators in the Gelfand–Tsetlin basis. It generalizes the well-known Gelfand–Tsetlin basis for the symmetric group (also known as Young's orthogonal form or Young–Yamanouchi basis).

We provide two applications of our result to quantum information. First, we show how to simplify semidefinite optimization problems over unitary-equivariant quantum channels by performing a symmetry reduction. Second, we derive an efficient quantum circuit for implementing the optimal port-based quantum teleportation protocol, exponentially improving the known trivial construction. As a consequence, this also exponentially improves the known lower bound for the amount of entanglement needed to implement unitaries non-locally.

Both applications require a generalization of quantum Schur transform to tensors of mixed unitary symmetry. We develop an efficient quantum circuit for this mixed quantum Schur transform and provide a matrix product state representation of its basis vectors. For constant local dimension, this yields an efficient classical algorithm for computing any entry of the mixed quantum Schur transform unitary.

Contents

1. Introduction 2
1.1. Background 2
1.2. Historical context 3
1.3. Strategy 4
1.4. Summary of our results 4
1.5. Related work 5
2. Preliminaries 6
2.1. Young diagrams and tableaux 6
2.2. Bratteli diagram and Gelfand–Tsetlin basis 7
2.3. Gelfand–Tsetlin patterns 9
2.4. Mixed Young diagrams and tableaux 9
3. Representation theory of partially transposed permutation matrix algebra 11
3.1. Walled Brauer algebra $\mathcal{B}^d_{p,q}$ 11
3.2. Matrix algebra $\mathcal{A}^d_{p,q}$ of partially transposed permutations 12
3.3. Representation theory of the algebra $\mathcal{A}^d_{p,q}$ 12
4. Mixed quantum Schur transform 15
4.1. Mixed Schur–Weyl duality 15
4.2. Mixed Schur transform 16
4.3. MPS representation of mixed Schur basis vectors 17
4.4. Mixed Schur transform achieves the Gelfand–Tsetlin basis 18
4.5. Quantum circuit for mixed Schur transform 19
4.6. Quantum circuit for Clebsch–Gordan transforms 19
5. Unitary-equivariant SDPs 22
5.1. Unitary-equivariant quantum channels 22
5.2. Full trace 23
5.3. Partial trace 24
5.4. Unitary-equivariant SDPs 25
6. Efficient quantum circuit for optimal port-based teleportation 27
6.1. Naimark's dilation 28
6.2. Efficient quantum circuit for the pretty good measurement 30

[1]Institute for Logic, Language, and Computation, University of Amsterdam and QuSoft, Amsterdam, The Netherlands
[2]Korteweg-de Vries Institute for Mathematics and Institute for Theoretical Physics, University of Amsterdam, The Netherlands
E-mail addresses: dmitry.grinko@cwi.nl, adam.burchardt@cwi.nl, marozols@gmail.com.