

2021–
2025

Gate-based counterdiabatic driving with complexity guarantees

Dyon van Vreumingen[1, 2]

[1] *Institute of Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands*
[2] *QuSoft, Centrum Wiskunde & Informatica (CWI), Science Park 123, 1098 XG Amsterdam, The Netherlands*

(Dated: October 10, 2024)

We propose a general, fully gate-based quantum algorithm for counterdiabatic driving. The algorithm does not depend on heuristics as in previous variational methods, and exploits regularisation of the adiabatic gauge potential to suppress only the transitions from the eigenstate of interest. This allows for a rigorous quantum gate complexity upper bound in terms of the minimum gap Δ around this eigenstate. We find that, in the worst case, the algorithm requires at most $O(\Delta^{-(3+o(1))}\epsilon^{-(1+o(1))})$ quantum gates to achieve a target state fidelity of at least $1-\epsilon^2$, where Δ is the minimum spectral gap. In certain cases, the gap dependence can be improved to quadratic.

arXiv:2406.08064v3 [quant-ph] 9 Oct 2024

I. INTRODUCTION

Adiabatic quantum computing (AQC) is a paradigm of quantum computation that leverages the principles of adiabatic processes in quantum mechanics to solve computational problems [1]. AQC relies on the adiabatic theorem, which states that a quantum system approximately remains in its instantaneous eigenstate if a given time-dependent hamiltonian that governs its energy levels is changed slowly enough [2]. In this sense, AQC is a conceptually straightforward way to prepare ground states or other eigenstates of complex systems, by starting from an eigenstate of a simple hamiltonian (e.g. a product state) and slowly time evolving towards the complex hamiltonian.

Contrary to AQC, counterdiabatic driving (CD) aims to prepare target states through a fast time evolution, actively suppressing excitations that would typically accompany such rapid driving. Traditional adiabatic processes require slow evolution to maintain the system in its instantaneous eigenstate, which can be impractically long. CD involves adding auxiliary, non-adiabatic control fields to counteract the diabatic transitions, thereby simulating the effect of a slow adiabatic process in a (much) shorter time [3]. It is a specific method within a broader class of techniques known as "shortcuts to adiabaticity" [4], designed to achieve the same goal of mimicking adiabatic evolution on a small time scale. Its original formulation is due to Demirplak & Rice [5–7]; in this formalism, the central object is the adiabatic gauge potential (AGP), which is an auxiliary field that is added to the system hamiltonian and cancels what is akin to a Coriolis force in the rotating system eigenbasis, thereby supressing all excitations. The issue however is that determining an AGP exactly for a given system is computationally difficult – possibly as difficult, if not more, as preparing the eigenstate of interest – and analytical expressions are only known for very specific systems [3]. This is due to the fact that, typically, the AGP is highly nonlocal and contains high-rank (two-body, three-body etc.) interactions [8–11]; moreover, if the spectrum of the system is gapless, the AGP does not exist [12].

For this reason, much effort has been focussed on computational methods that approximate the AGP. In this context, a pivotal role has been played by variational methods [13]; these methods assume a parametrised ansatz that is optimised with respect to an action function that approaches zero as the ansatz approaches the exact AGP. This approach has enabled approximate CD in a large number of different contexts, leading to improved fidelities as compared to standard AQC [14–20]. Among these variational methods, gate-based (digitised) approaches have been proposed as well [21], where the dynamics is trotterised for implementation on a gate-based quantum computer. However, the limit to the variational approach lies in the fact that evaluating the action function itself becomes harder when higher-rank (two-body, three-body etc.) interactions are taken into account in the ansatz; therefore only ansätze with low-rank interactions are computationally feasible for variational methods.

Further improvements to this scheme were made through the invention of a nested commutator ansatz to the AGP [22] and subsequent Krylov space methods to solve for the optimal coefficients in this ansatz [23, 24]. Here, Krylov spaces are subspaces of operator space that are specifically chosen such that the coefficients can be found by solving a linear equation, which avoids optimisation heuristics with unpredictable running times. At the same time, this approach sheds more light on the computational complexity of computing the AGP. Numerical experiments [23] suggest that the relevant search space for weakly interacting and integrable systems is a small subspace in operator space, so that the AGP is easily approximated with an ansatz containing few coefficients; on the other hand, for strongly interacting, chaotic systems, this tends not to be the case, and the whole operator space may need to be explored. In this sense, the complexity of CD is tightly connected to quantum Krylov complexity [25–30]. More recently, interesting suggestions have been put forward to apply tensor network approaches for approximating the