


Continuous cavity-QED with an atomic beam

Francesca Famà,[1] Sheng Zhou,[1] Benedikt Heizenreder,[1] Mikkel Tang,[1,2] Shayne Bennetts,[1] Simon B. Jäger,[3] Stefan A. Schäffer,[1,2] and Florian Schreck[1,4,*]

[1] *Van der Waals-Zeeman Institute, Institute of Physics, University of Amsterdam, Science Park 904, 1098XH Amsterdam, The Netherlands*
[2] *Niels Bohr Institute, University of Copenhagen, Blegdamsvej 17, 2100 Copenhagen, Denmark*
[3] *Physics Department and Research Center OPTIMAS, University of Kaiserslautern-Landau, 67663 Kaiserslautern, Germany*
[4] *QuSoft, Science Park 123, 1098XG Amsterdam, The Netherlands*

(Dated: July 29, 2024)

Atoms coupled to cavities provide an exciting playground for the study of fundamental interactions of atoms mediated through a common channel. Many of the applications of cavity-QED and cold-atom experiments more broadly, suffer from limitations caused by the transient nature of an atomic loading cycle. The development of continuous operation schemes is necessary to push these systems to the next level of performance. Here we present a machine designed to produce a continuous flux of collimated atoms that traverse an optical cavity. The atom-light interaction is enhanced by a fast-decaying cavity optimal for studying phenomena where atomic properties dominate. We demonstrate the transition to a collective strong coupling regime heralded by a normal-mode splitting. We observe a second phase with a binary normal-mode splitting born from an offset in the mean velocity of the atoms. Inverting the atomic ensemble in the collective strong coupling regime, we measure continuous optical gain. This work sets the stage for studying threshold conditions for continuous collective phenomena, such as continuous superradiant lasing.

arXiv:2407.18668v1 [physics.atom-ph] 26 Jul 2024

I. INTRODUCTION

Developing techniques to realize the continuous operation of quantum systems based on cold atoms is a long-standing goal within the quantum community [1–7]. This requires that atoms are continuously loaded into the region of interest, as their typical trapping times are usually limited by heating and decoherence processes [8]. Consequently, one of the main challenges arises from the finite interaction time of the individual atoms and whether it fundamentally limits the coherence time of the quantum system [9]. Overcoming this challenge will result in the realization of continuous cold atom platforms, which will find broad applications in devices including quantum simulators and computers [10], but also ultra-precise quantum sensors and clocks [11–13].

A paradigmatic example of such a device is a continuous beam of cold atoms that traverses and interacts with an optical cavity, which can mediate interaction between the atoms. For cavity decay rates much larger than the other rates involved, the cavity field will have a short memory of phase information. This case is often referred to as the 'bad-cavity' regime, in which the cavity acts as a dissipative entity, providing a preferred vacuum channel to which all atoms couple [14]. Especially interesting is the case of a narrow, semi-forbidden atomic transition couples to the optical cavity. Such a coupled system might be used to realize the next generation of superradiant lasers and active atomic clocks that operate in continuous-wave mode [15, 16]. Devices working in the bad-cavity regime, can be extremely robust against environmental noise [17, 18]. Furthermore, they can overcome the issue of discontinuous phase evolution emerging from cyclic operation, seen, e.g., as the Dick effect, which limits the frequency stability of state-of-the-art atomic clocks [19, 20]. The phase-noise introduced through cycling significantly affects the time scaling by limiting the $1/\text{time}$ scaling of white phase noise to $1/\sqrt{\text{time}}$ at all operating time scales. On the fundamental side, such systems are formidable playgrounds for exploring many-body physics emerging from the interplay of continuous atom flow and long-range cavity-mediated atom-atom interactions [21]. Collective atom-cavity strong coupling has been observed in (quasi-)continuous systems for narrow transitions with ultracold atoms in a cavity [22–24], and in thermal atoms transiting a cavity mode, with broad atomic transitions coupled to the same mode [25]. However, it remains an open question whether the cavity and atoms can form steady-state coherent dressed states for a semiforbidden atomic transition, having to overcome the finite lifetimes of cavity photons and atoms. For an atomic beam, this is particularly challenging as the lifetime of the atoms in the cavity mode is determined by their finite transit time rather than their natural, narrow linewidth. The demonstration that coherent collective atom-cavity interactions can overcome these processes is an important step towards establishing such devices as future quantum technologies [26].

In this paper, we demonstrate the formation of dressed states of a thermal atomic beam that traverses an optical cavity, see Fig. 1(c). In particular, we report strong collective light-matter coupling of the narrow 7.5 kHz ${}^1S_0 \leftrightarrow {}^3P_1$ transition of ^{88}Sr to a bad cavity. The transit times of the atoms is the order of hundreds of nanoseconds, while

* continuousCollectiveGain@strontiumBEC.com