

Mutual information fluctuations and non-stabilizerness in random circuits

Arash Ahmadi,[1,*] Jonas Helsen,[2] Cagan Karaca,[1] and Eliska Greplova[1]

[1] *QuTech and Kavli Institute of Nanoscience, Delft University of Technology, Delft, the Netherlands*
[2] *QuSoft and CWI, Amsterdam, The Netherlands*

The emergence of quantum technologies has brought much attention to the characterization of quantum resources as well as the classical simulatability of quantum processes. Quantum resources, as quantified by non-stabilizeress, have in one theoretical approach been linked to a family of entropic, monotonic functions. In this work, we demonstrate both analytically and numerically a simple relationship between non-stabilizerness and information scrambling using the fluctuations of an entropy-based quantifier. Specifically, we find that the non-stabilizerness generated by a random quantum circuit is proportional to fluctuations of mutual information. Furthermore, we explore the role of non-stabilizerness in measurement-induced entanglement phase transitions. We find that the fluctuations of mutual information decrease with increasing non-stabilizerness yielding potentially easier identification of the transition point. Our work establishes a key connection between quantum resource theory, information scrambling and measurement-induced entanglement phase transitions.

arXiv:2408.03831v2 [quant-ph] 12 Aug 2024

I. INTRODUCTION

Since the formulation of quantum theory, entanglement has been known to be one of the unique signatures of quantum theory [1]. While entanglement is one of the key features of distinguishing the quantum world from the classical, it is known that entanglement alone is insufficient for having an advantage from the *computational* point of view. In particular, the stabilizer states are a set of states that can be highly entangled and at the same time can be simulated efficiently on classical computers [2–7].

The missing feature for quantum computation to have an advantage over classical computation is a concept known as non-stabilizerness or *magic* [8]. In quantum circuit language, in the fault-tolerant regime, the gates that are known to be "cheap" gates [8] are Clifford gates, and magic is injected into the state by adding non-Clifford gates to the circuit. It is also known that the Clifford operations could have easier implementation both at the experimental level and for quantum error correction [2, 4, 9, 10]. This feature makes magic the resource for quantum computation.

Various measures for non-stabilizerness, or magic, have been proposed. Some of the notable ones are magical cross-entropy, mana [8], robustness of magic [11, 12], stabilizer entropies (SE) [13] and others [14–18]. Due to the favourable scaling of the stabilizer entropies these measures received much attention from the community to study the properties of non-stabilizerness of quantum systems [19–25]. Additionally, there has been a significant effort to scale up the computability of stabilizer entropies [26–30].

Diverse studies confirmed there exists a relationship between non-stabilizerness and information scrambling [13, 16, 31]. Information scrambling describes the spread of local information in a generic quantum system [32]. Information scrambling has been shown to be one of the most powerful measures for various quantum properties of quantum systems, from black holes [33–36] to many-body quantum systems [37] to quantum circuits [31, 38]. There are a number of known methods for measuring information scrambling. One of the well-known approaches is based on the measurement of correlator functions, namely Out-of-Time Ordered Correlators (OTOCs) [33, 36, 38–41]. Another approach for studying information scrambling are entropy-based measures [32, 42–48].

In this paper, we describe numerically and analytically a general relationship between non-stabilizerness and *fluctuations* of entropy-based measures of information scrambling. A specific instance of this relation has been observed earlier by connecting fluctuations of OTOCs to non-stabilizerness [13, 31]. In the present work, we generalize this behaviour to fluctuations of generic mutual information measures of disjoint regions of the quantum circuit on t-doped circuits. The observed link between non-stabilizerness and mutual information creates a bridge to the theory of phase transitions in measurement-induced random quantum circuits, where mutual information plays a crucial role [49–81]. Here we show that non-stabilizerness is an ingredient in reducing the spread of entanglement phase transition in random quantum circuits and also reduces the fluctuations of the measured mutual information.

II. RANDOM CIRCUITS AND MUTUAL INFORMATION

A. Definitions and notation

The structure of the T-doped circuits that we use in this study consists of blocks of random Clifford operations followed by single T-gates on random qubits in the circuit. The blocks of random Clifford consist of single Clifford gates drawn randomly from the set $\{I, X, Y, Z, H, S\}$ followed by three CNOT gates on two

* a.ahmadi-1@tudelft.nl