

2022–
2024

arXiv:2408.07067v2 [quant-ph] 19 Sep 2024

Asymptotic quantification of entanglement with a single copy

Ludovico Lami,[1,2,3,*] Mario Berta,[4,†] and Bartosz Regula[5,‡]

[1]*QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*

[2]*Korteweg–de Vries Institute for Mathematics, University of Amsterdam, Science Park 105-107, 1098 XG Amsterdam, the Netherlands*

[3]*Institute for Theoretical Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands*

[4]*Institute for Quantum Information, RWTH Aachen University, Aachen, Germany*

[5]*Mathematical Quantum Information RIKEN Hakubi Research Team, RIKEN Cluster for Pioneering Research (CPR) and RIKEN Center for Quantum Computing (RQC), Wako, Saitama 351-0198, Japan*

Despite the central importance of quantum entanglement in fueling many quantum technologies, the understanding of the optimal ways to exploit it is still beyond our reach, and even measuring entanglement in an operationally meaningful way is prohibitively difficult. This is due to the need to precisely characterise many-copy, asymptotic protocols for entanglement processing. Here we overcome these issues by introducing a new way of benchmarking the fundamental protocol of entanglement distillation (purification), where instead of measuring its asymptotic yield, we focus on the best achievable error. We connect this formulation of the task with an information-theoretic problem in composite quantum hypothesis testing known as generalised Sanov's theorem. By solving the latter problem — which had no previously known solution even in classical information theory — we thus compute the optimal asymptotic error exponent of entanglement distillation. We show this asymptotic solution to be given by the *reverse relative entropy of entanglement*, a single-letter quantity that can be evaluated using only a single copy of a quantum state, which is a unique feature among operational measures of entanglement. Altogether, we thus demonstrate a measure of entanglement that admits a direct operational interpretation as the optimal asymptotic rate of an important entanglement manipulation protocol while enjoying an exact, single-letter formula.

I. INTRODUCTION

The phenomenon of quantum entanglement is one of the most important resources that underlie the potential of quantum technologies to provide advantages in information processing and computation [1–6]. And yet its precise characterisation remains very little understood. Take for instance the key task of entanglement distillation, concerned with purifying noisy entanglement into a pure, maximally entangled form [7–9]. This task is a vital primitive that underlies practical protocols that utilise entanglement, as the pure form of entanglement is required for most quantum computation and communication schemes. Despite this clear significance, and despite being one of the very first operational protocols ever studied in quantum information theory, we still do not have a computable formula for how much entanglement can be distilled from a given quantum state — even deciding whether any entanglement whatsoever can be extracted is an unsolved problem in general [10, 11]. Similar problems affect other entanglement processing tasks, and exact solutions generally exist only in few special cases.

* ludovico.lami@gmail.com
† berta@physik.rwth-aachen.de
‡ bartosz.regula@gmail.com