


Phonon-mediated quantum gates in trapped ions coupled to an ultracold atomic gas

Lorenzo Oghittu[1,2], Arghavan Safavi-Naini[1,2], Antonio Negretti[3] and Rene Gerritsma[1,4]

[1] *Institute for Theoretical Physics, Institute of Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands,*
[2] *QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands,*
[3] *Zentrum für Optische Quantentechnologien, Universität Hamburg, Luruper Chaussee 149, 22761 Hamburg, Germany,*
[4] *Van der Waals-Zeeman Institute, Institute of Physics, University of Amsterdam, 1098 XH Amsterdam, the Netherlands*

(Dated: December 9, 2024)

We study the dynamics of phonon-mediated qubit-qubit interactions between trapped ions in the presence of an ultracold atomic gas. By deriving and solving a master equation to describe the combined system, we show that the presence of the atoms causes the quantum gate quality to reduce because of motional decoherence. On the other hand, we calculate that the gas may be used to keep the ion crystal cold in the presence of external heating due to electric field noise. We show that tuning the atom-ion scattering length allows one to tune the cooling rate of the ions and would make it possible to temporarily reduce the effects of the gas during a quantum gate while keeping the ions cold over long timescales. In this way, the trapped ion quantum computer may be buffer gas cooled. The system may also be used for quantum-enhanced measurements of the atom-ion interactions or properties of the atomic bath.

arXiv:2409.03693v2 [quant-ph] 6 Dec 2024

I. INTRODUCTION

Over the last decade, rapid progress has occurred in controlling laboratory systems that explore the quantum dynamics of mixtures of ultracold atoms and trapped ions [1–10]. These systems offer new opportunities in studying quantum impurity physics, ultracold collisions and quantum chemistry [11–17]. On the theoretical side, efforts have been made to study and characterize ionic polarons [18–20] and bath-mediated interactions between charged impurities [21–23]. Moreover, systems involving ionic impurities in ultracold gases have been proposed as quantum simulators and sensors [24–27]. Combining atom-ion systems with precise quantum control of individual trapped ions opens the possibility to study the decoherence of non-classical states of ion qubits and motion [28, 29] as well as phonon-mediated interactions between ion qubit states [30–32] inside a bath of ultracold atoms. These mixed atom-ion systems may find applications in quantum-enhanced sensing of atom-ion interactions [17] and the properties of the atomic bath [33]. Furthermore, the ultracold atomic gas may be used to cool the trapped ions, with recent results showing that buffer gas cooling even outperforms laser (Doppler) cooling [9, 34]. Finally, the observation of atom-ion Feshbach resonances has greatly enhanced the possibilities of atom-ion quantum systems [35]. In particular, these resonances allow for controlling the atom-ion interactions by simple tuning of an external magnetic field, in analogy to the case of Feshbach resonances between neutral atoms [36].

In this paper, we consider the case where ionic qubits are coupled via laser-driven phonon-mediated interactions and are coupled to a quantum bath of ultracold atoms. We study how the atoms change the dynamics of this system and affect the quality of quantum gates between the qubits. We find that the main source of error occurs due to decoherence of the ion motion in phase space during the gate. We consider the benefits of the atomic bath to keep the ions cool in the presence of an external heating source due to electric field noise. We study the role of the atom-ion s-wave scattering length, which may be tuned using Feshbach resonances. We find that it is possible to switch the interactions in such a way that either the adverse effects of the bath on the quantum gate or the negative effects of heating may be mitigated. These results show that it is possible to buffer gas cool the trapped ion quantum system [37], while maintaining the quality of quantum operations. Moreover, our model involves a setup that is experimentally accessible to several groups working on hybrid atom-ion systems, and therefore paves the way for the study of combined systems of ultracold atoms and quantum gates.




FIG. 1. Scheme of the system. The three ions are described as a one dimensional chain whose motion is decomposed into three normal modes of oscillation (dashed lines). Ions 1 and 3 are considered as two-level spin systems. Ion 2 is immersed in a bath of ultracold atoms (red particles) which are assumed to be effectively free

. A thermal bath at room temperature (wavy line) is coupled to the center-of-mass motion of the ion chain.

The paper is organized as follows: in Sec. II we describe the Hamiltonian of the system, which is used in Sec. III