

2022–
2024

arXiv:2501.12447v4 [quant-ph] 11 Dec 2025

Tight relations and equivalences between smooth relative entropies

Bartosz Regula,[1,*] Ludovico Lami,[2,3,4,5,†] and Nilanjana Datta[6,‡]

[1]*Mathematical Quantum Information RIKEN Hakubi Research Team, RIKEN Pioneering Research Institute (PRI) and RIKEN Center for Quantum Computing (RQC), Wako, Saitama 351-0198, Japan*
[2]*Scuola Normale Superiore, Piazza dei Cavalieri 7, 56126 Pisa, Italy*
[3]*QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*
[4]*Korteweg–de Vries Institute for Mathematics, University of Amsterdam, Science Park 105-107, 1098 XG Amsterdam, the Netherlands*
[5]*Institute for Theoretical Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands*
[6]*Department of Applied Mathematics and Theoretical Physics, Centre for Mathematical Sciences, University of Cambridge, Cambridge CB3 0WA, United Kingdom*

The precise one-shot characterisation of operational tasks in classical and quantum information theory relies on different forms of smooth entropic quantities. A particularly important connection is between the hypothesis testing relative entropy and the smoothed max-relative entropy, which together govern many operational settings. We first strengthen this connection into a type of equivalence: we show that the hypothesis testing relative entropy is equivalent to a variant of the smooth max-relative entropy based on the information spectrum divergence, which can be alternatively understood as a measured smooth max-relative entropy. Furthermore, we improve a fundamental lemma due to Datta and Renner that connects the different variants of the smoothed max-relative entropy, introducing a modified proof technique based on matrix geometric means and a tightened gentle measurement lemma. We use the unveiled connections and tools to strictly improve on previously known one-shot bounds and duality relations between the smooth max-relative entropy and the hypothesis testing relative entropy, establishing provably tight bounds between them. We use these results to refine other divergence inequalities, in particular sharpening bounds that connect the max-relative entropy with Rényi divergences.

CONTENTS

I. Introduction 2

II. Summary of main results 3

III. Smooth divergences 4
A. Hypothesis testing relative entropy and max-relative entropy 4
B. Modified max-relative entropy $\widetilde{D}^{\varepsilon}_{\max}$ 7

IV. On normalised and subnormalised smoothing 7

V. Equivalences 10
A. Classical $\widetilde{D}^{\varepsilon}_{\max}$ and $D^{\varepsilon}_{\max}$ 10
B. $\widetilde{D}^{\varepsilon}_{\max}$ as measured $D^{\varepsilon}_{\max}$ 10
C. D^{ε}_{H} as optimised $\widetilde{D}^{\varepsilon}_{\max}$ 12

* bartosz.regula@gmail.com
† ludovico.lami@gmail.com
‡ n.datta@damtp.cam.ac.uk