

2023–
2024



2022–
2024



arXiv:2503.21479v2 [quant-ph] 9 Oct 2025

Quantum umlaut information

Filippo Girardi,[1,2,3,*] Aadil Oufkir,[4,†] Bartosz Regula,[5] Marco Tomamichel,[6,7] Mario Berta,[4] and Ludovico Lami[1,2,3]

[1]*Scuola Normale Superiore, Piazza dei Cavalieri 7, 56126 Pisa, Italy*

[2]*QuSoft, Science Park 123, 1098 XG Amsterdam, The Netherlands*

[3]*Korteweg–de Vries Institute for Mathematics, University of Amsterdam, Science Park 105-107, 1098 XG Amsterdam, The Netherlands*

[4]*Institute for Quantum Information, RWTH Aachen University, Germany*

[5]*Mathematical Quantum Information RIKEN Hakubi Research Team, RIKEN Pioneering Research Institute (PRI) and RIKEN Center for Quantum Computing (RQC), Wako, Saitama 351-0198, Japan*

[6]*Centre for Quantum Technologies, National University of Singapore, Singapore*

[7]*Department of Electrical and Computer Engineering, National University of Singapore, Singapore*

We study the quantum umlaut information, a correlation measure defined for bipartite quantum states ρ_{AB} as a reversed variant of the quantum mutual information: $U(A;B)_\rho = \min_{\sigma_B} D(\rho_A \otimes \sigma_B \| \rho_{AB})$ in terms of the quantum relative entropy D. As in the classical case [Girardi et al., arXiv:2503.18910], this definition allows for a closed-form expression and has an operational interpretation as the asymptotic error exponent in the hypothesis testing task of deciding whether a given bipartite state is product or not. We generalise the umlaut information to quantum channels, where it also extends the notion of 'oveloh information' [Nuradha et al., J. Phys. A 58, 165304 (2025)]. We prove that channel umlaut information is additive for classical-quantum channels, while we observe additivity violations for fully quantum channels. Inspired by recent results in entanglement theory, we then show as our main result that the regularised umlaut information constitutes a fundamental measure of the *quality* of classical information transmission over a quantum channel — as opposed to the capacity, which quantifies the *quantity* of information that can be sent. This interpretation applies to coding assisted by activated non-signalling correlations, and the channel umlaut information is in general larger than the corresponding expression for unassisted communication as obtained by Dalai for the classical-quantum case [IEEE Trans. Inf. Theory 59, 8027 (2013)]. Combined with prior works on non-signalling–assisted zero-error channel capacities, our findings imply a dichotomy between the settings of zero-rate error exponents and zero-error communication. While our results are single-letter only for classical-quantum channels, we also give a single-letter bound for fully quantum channels in terms of the 'geometric' version of umlaut information.

I. INTRODUCTION

A. Motivation

The quantification of the information transmission capabilities of quantum channels is a fundamental problem in quantum information theory, with prominent applications in quantum technologies. Most of the literature focused on the question of computing the asymptotic *capacities* of channels, i.e. the maximum amount of information (either classical or quantum) that can be

* filippo.girardi@sns.it
† oufkir@physik.rwth-aachen.de