

2023–
2025



2021–
2026



arXiv:2509.09813v1 [quant-ph] 11 Sep 2025

Nearly optimal algorithms to learn sparse quantum Hamiltonians in physically motivated distances

Amira Abbas
Google Quantum AI

Nunzia Cerrato
Scuola Normale Superiore

Francisco Escudero Gutiérrez
Centrum Wiskunde & Informatica (CWI)
and QuSoft

Dmitry Grinko
ILLC, University of Amsterdam
and QuSoft

Francesco Anna Mele
NEST, Scuola Normale Superiore
and Istituto Nanoscienze

Pulkit Sinha
Institute for Quantum Computing,
University of Waterloo

Abstract

We study the problem of learning Hamiltonians H that are s-sparse in the Pauli basis, given access to their time-evolution operators. Although Hamiltonian learning has been extensively investigated, two issues recur in much of the existing literature: the absence of lower bounds establishing optimality and the use of mathematically convenient but physically opaque error measures.

We address both challenges by introducing two physically motivated notions of distance between Hamiltonians and designing a nearly optimal algorithm with respect to one of these metrics. The first, the *time-constrained distance*, quantifies distinguishability through dynamical evolution up to a bounded time. The second, the *temperature-constrained distance*, captures distinguishability through thermal states at bounded inverse temperatures.

We show that s-sparse Hamiltonians with bounded operator norm can be learned under both distances using only $\widetilde{O}(s\log(1/\varepsilon))$ experiments and $\widetilde{O}(s^2/\varepsilon)$ total evolution time. For the time-constrained distance, we further establish lower bounds of $\Omega((s/n)\log(1/\varepsilon)+s)$ experiments and $\Omega(\sqrt{s}/\varepsilon)$ total evolution time, demonstrating near-optimality in the number of experiments.

As an intermediate result, we obtain an algorithm that learns every Pauli coefficient of s-sparse Hamiltonians up to error ε in $\widetilde{O}(s\log(1/\varepsilon))$ experiments and $\widetilde{O}(s/\varepsilon)$ total evolution time, improving upon several recent results.

The source of this improvement is a new *isolation technique*, inspired by the Valiant-Vazirani theorem (STOC'85), which shows that *NP is as easy as detecting unique solutions*. This isolation technique allows us to query the time evolution of a single Pauli coefficient of a sparse Hamiltonian—even when the Pauli support of the Hamiltonian is unknown—ultimately enabling us to recover the Pauli support itself.